

**CNDP1 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP9166b****Specification**

---

**CNDP1 Antibody (C-term) - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB, IHC-P, FC,E        |
| Primary Accession | <a href="#">Q96KN2</a> |
| Reactivity        | Human                  |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |
| Isotype           | Rabbit IgG             |
| Calculated MW     | 56692                  |
| Antigen Region    | 415-443                |

**CNDP1 Antibody (C-term) - Additional Information****Gene ID** 84735**Other Names**

Beta-Ala-His dipeptidase, CNDP dipeptidase 1, Carnosine dipeptidase 1, Glutamate carboxypeptidase-like protein 2, Serum carnosinase, CNDP1, CN1, CPGL2

**Target/Specificity**

This CNDP1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 415-443 amino acids from the C-terminal region of human CNDP1.

**Dilution**

WB~~1:1000  
IHC-P~~1:50~100  
FC~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CNDP1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**CNDP1 Antibody (C-term) - Protein Information****Name** CNDP1 ([HGNC:20675](#))

**Synonyms** CN1, CPGL2

**Function** Catalyzes the peptide bond hydrolysis in Xaa-His dipeptides, displaying the highest activity toward carnosine (beta-alanyl-L- histidine) and anserine (beta-alanyl-3-methyl-histidine).

**Cellular Location**

Secreted.


**Tissue Location**

Found in serum and adult nervous central system. Absent in serum from patients with homocarnosinosis


**CNDP1 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**CNDP1 Antibody (C-term) - Images**

Western blot analysis of CNDP1 Antibody (C-term) (Cat. #AP9166b) in CEM cell line lysates (35ug/lane). CNDP1 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human brain tissue reacted with CNDP1 Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



CNDP1 Antibody (C-term) (Cat. #AP9166b) flow cytometry analysis of CEM cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

### CNDP1 Antibody (C-term) - Background

CNDP1 encodes a member of the M20 metalloprotease family. The encoded protein is specifically expressed in the brain, is a homodimeric dipeptidase which was identified as human carnosinase. This protein contains trinucleotide (CTG) repeat length polymorphism in the coding region.

### CNDP1 Antibody (C-term) - References

Craig, D.W., et al., *Diabet. Med.* 26 (11), 1090-1098 (2009)  
Mooyaart, A.L., et al., *Diabetes Res. Clin. Pract.* 85 (3), 272-278 (2009)