

ACCN1 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP9213c**Specification**

ACCN1 Antibody (Center) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	Q16515
Other Accession	Q62962 , Q925H0
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	57709
Antigen Region	120-148

ACCN1 Antibody (Center) - Additional Information**Gene ID 40****Other Names**

Acid-sensing ion channel 2, ASIC2, Amiloride-sensitive brain sodium channel, Amiloride-sensitive cation channel 1, neuronal, Amiloride-sensitive cation channel neuronal 1, Brain sodium channel 1, BNC1, BNaC1, Mammalian degenerin homolog, ASIC2, ACCN, ACCN1, BNAC1, MDEG

Target/Specificity

This ACCN1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 120-148 amino acids from the Central region of human ACCN1.

Dilution

WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ACCN1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

ACCN1 Antibody (Center) - Protein Information

Name ASIC2

Synonyms ACCN, ACCN1, BNAC1, MDEG

Function Cation channel with high affinity for sodium, which is gated by extracellular protons and inhibited by the diuretic amiloride. Also permeable for Li(+) and K(+). Generates a biphasic current with a fast inactivating and a slow sustained phase. Heteromeric channel assembly seems to modulate.

Cellular Location

Cell membrane; Multi-pass membrane protein. Note=Localized at the plasma membrane of neurons, in the soma and punctated peripheral processes.

Tissue Location


Brain and spinal cord. Isoform 1 is also detected in testis, liver, colon and ovary.

ACCN1 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


ACCN1 Antibody (Center) - Images



Western blot analysis of ACCN1 Antibody (Center) (Cat. #AP9213c) in NCI-H460 cell line lysates (35ug/lane). ACCN1 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human brain tissue reacted with ACCN1 Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



ACCN1 Antibody (Center) (Cat. #AP9213c) flow cytometric analysis of WiDr cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

ACCN1 Antibody (Center) - Background

ACCN1 encodes a member of the degenerin/epithelial sodium channel (DEG/ENaC) superfamily. The members of this family are amiloride-sensitive sodium channels that contain intracellular N and C termini, 2 hydrophobic transmembrane regions, and a large extracellular loop, which has many cysteine residues with conserved spacing. The member encoded by this protein may play a role in neurotransmission. In addition, a heteromeric association between this member and ACCN3 (variant 1) has been observed to co-assemble into proton-gated channels sensitive to gadolinium.

ACCN1 Antibody (Center) - References

Bashari, E., et al., Am. J. Physiol., Cell Physiol. 296 (2), C372-C384 (2009)
 Chai, S., et al., J. Biol. Chem. 282 (31), 22668-22677 (2007)