10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

FMOD Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP9243b

Specification

FMOD Antibody (C-term) - Product Information

Application	**WB, IHC-P, FC,E**
Primary Accession	Q06828
Other Accession	P13605
Reactivity	**Human, Mouse**
Predicted	**Bovine**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**43179**
Antigen Region	**337-364**

FMOD Antibody (C-term) - Additional Information

Gene ID 2331

Other Names
Fibromodulin, FM, Collagen-binding 59 kDa protein, Keratan sulfate proteoglycan fibromodulin, KSPG fibromodulin, FMOD, FM, SLRR2E

Target/Specificity
This FMOD antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 337-364 amino acids from the C-terminal region of human FMOD.

Dilution
WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
FMOD Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

FMOD Antibody (C-term) - Protein Information

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Name FMOD

Synonyms FM, SLRR2E

Function Affects the rate of fibrils formation. May have a primary role in collagen fibrillogenesis (By similarity).


Cellular Location
Secreted, extracellular space, extracellular matrix

FMOD Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

FMOD Antibody (C-term) - Images




Western blot analysis of FMOD Antibody (C-term) (Cat. #AP9243b) in mouse cerebellum tissue lysates (35ug/lane). FMOD (arrow) was detected using the purified Pab.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



Formalin-fixed and paraffin-embedded human brain tissue reacted with FMOD Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



FMOD Antibody (C-term) (Cat. #AP9243b) flow cytometric analysis of Hela cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

FMOD Antibody (C-term) - Background

Fibromodulin is a member of a family of small interstitial proteoglycans, containing a central region composed of leucine-rich repeats with 4 keratan sulfate chains flanked by disulfide-bonded terminal domains. It may participate in the assembly of the extracellular matrix as it interacts with type I and type II collagen fibrils and inhibits fibrillogenesis in vitro. It may also regulate TGF-beta activities by sequestering TGF-beta into the extracellular matrix.

FMOD Antibody (C-term) - References

Ehret,G.B., et.al, Eur. J. Hum. Genet. 17 (12), 1650-1657 (2009)
Tillgren,V., et.al, J. Biol. Chem. 284 (42), 28543-28553 (2009)