

DIRAS1 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP9402c

Specification

DIRAS1 Antibody (Center) - Product Information

Application WB, IHC-P, FC,E

Primary Accession
Reactivity
Human
Host
Clonality
Polyclonal
Isotype
Calculated MW
Antigen Region
O95057
Human
Rabbit
Rabbit
Polyclonal
Rabbit IgG
Ca2329
Antigen Region
109-136

DIRAS1 Antibody (Center) - Additional Information

Gene ID 148252

Other Names

GTP-binding protein Di-Ras1, Distinct subgroup of the Ras family member 1, Ras-related inhibitor of cell growth, Rig, Small GTP-binding tumor suppressor 1, DIRAS1, GBTS1, RIG

Target/Specificity

This DIRAS1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 109-136 amino acids from the Central region of human DIRAS1.

Dilution

WB~~1:1000 IHC-P~~1:50~100 FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DIRAS1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

DIRAS1 Antibody (Center) - Protein Information

Name DIRAS1





Synonyms GBTS1, RIG

Function Displays low GTPase activity and exists predominantly in the GTP-bound form.

Cellular Location

Cell membrane; Lipid-anchor; Cytoplasmic side

Tissue Location


Highly expressed in heart and brain.

DIRAS1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

DIRAS1 Antibody (Center) - Images




Western blot analysis of DIRAS1 Antibody (Center) (Cat. #AP9402c) in CEM cell line lysates (35ug/lane). DIRAS1 (arrow) was detected using the purified Pab.





Formalin-fixed and paraffin-embedded human brain tissue reacted with DIRAS1 Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



DIRAS1 Antibody (Center) (Cat. #AP9402c) flow cytometry analysis of MDA-MB231 cells (bottom histogram) compared to a negative control cell (top histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

DIRAS1 Antibody (Center) - Background

DIRAS1 belongs to a distinct branch of the functionally diverse Ras (see HRAS; MIM 190020) superfamily of monomeric GTPases.

DIRAS1 Antibody (Center) - References

Kimura, K., et al. Genome Res. 16(1):55-65(2006) Kontani, K., et al. J. Biol. Chem. 277(43):41070-41078(2002) Ellis, C.A., et al. Proc. Natl. Acad. Sci. U.S.A. 99(15):9876-9881(2002)