

FBXL5 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP9409a**Specification**

FBXL5 Antibody (N-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	Q9UKA1
Other Accession	Q8C2S5 , A2VE78
Reactivity	Human, Mouse
Predicted	Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	86-115

FBXL5 Antibody (N-term) - Additional Information**Gene ID** 26234**Other Names**

F-box/LRR-repeat protein 5, F-box and leucine-rich repeat protein 5, F-box protein FBL4/FBL5, p45SKP2-like protein, FBXL5, FBL4, FBL5, FLR1

Target/Specificity

This FBXL5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 86-115 amino acids from the N-terminal region of human FBXL5.

Dilution

WB~~1:1000
IHC-P~~1:100
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FBXL5 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

FBXL5 Antibody (N-term) - Protein Information**Name** FBXL5

Synonyms FBL4, FBL5, FLR1

Function Component of some SCF (SKP1-cullin-F-box) protein ligase complex that plays a central role in iron homeostasis by promoting the ubiquitination and subsequent degradation of IREB2/IRP2 (PubMed:[19762596](#), PubMed:[19762597](#)). The C-terminal domain of FBXL5 contains a redox-sensitive [2Fe-2S] cluster that, upon oxidation, promotes binding to IRP2 to effect its oxygen-dependent degradation (PubMed:[32126207](#)). Under iron deficiency conditions, the N-terminal hemerythrin-like (Hr) region, which contains a diiron metal center, cannot bind iron and undergoes conformational changes that destabilize the FBXL5 protein and cause its ubiquitination and degradation (PubMed:[19762596](#), PubMed:[19762597](#)). When intracellular iron levels start rising, the Hr region is stabilized (PubMed:[19762596](#), PubMed:[19762597](#)). Additional increases in iron levels facilitate the assembly and incorporation of a redox active [2Fe-2S] cluster in the C-terminal domain (PubMed:[32126207](#)). Only when oxygen level is high enough to maintain the cluster in its oxidized state can FBXL5 recruit IRP2 as a substrate for polyubiquitination and degradation (PubMed:[32126207](#)). Promotes ubiquitination and subsequent degradation of the dynactin complex component DCTN1 (PubMed:[17532294](#)). Within the nucleus, promotes the ubiquitination of SNAI1; preventing its interaction with DNA and promoting its degradation (PubMed:[24157836](#)). Negatively regulates DNA damage response by mediating the ubiquitin-proteasome degradation of the DNA repair protein NABP2 (PubMed:[25249620](#)).


Cellular Location

Cytoplasm, perinuclear region. Nucleus


FBXL5 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FBXL5 Antibody (N-term) - Images


All lanes : Anti-FBXL5 Antibody (N-term) at 1:1000 dilution Lane 1: Hela whole cell lysate Lane 2: U-87 MG whole cell lysate Lane 3: HepG2 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 79 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



Immunohistochemical analysis of AP9409A on paraffin-embedded Human kidney tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



FBXL5 Antibody (N-term) (Cat. #AP9409a) flow cytometric analysis of HepG2 cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Western blot analysis of FBXL5 Antibody (N-term) (Cat. #AP9409a) in mouse heart tissue lysates (35ug/lane). FBXL5 (arrow) was detected using the purified Pab.

FBXL5 Antibody (N-term) - References

- Salahudeen, A.A., et al. Science 326(5953):722-726(2009)
Vashisht, A.A., et al. Science 326(5953):718-721(2009)
Zhang, N., et al. Biochem. Biophys. Res. Commun. 359(1):34-39(2007)
Ilyin, G.P., et al. Genomics 67(1):40-47(2000)
Winston, J.T., et al. Curr. Biol. 9(20):1180-1182(1999)
Cenciarelli, C., et al. Curr. Biol. 9(20):1177-1179(1999)