

MAP4K3 Antibody
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP9860A**Specification**

MAP4K3 Antibody - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	Q8IVH8
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	101316

MAP4K3 Antibody - Additional Information**Gene ID** 8491**Other Names**

Mitogen-activated protein kinase kinase kinase kinase 3, Germinal center kinase-related protein kinase, GLK, MAPK/ERK kinase kinase kinase 3, MEK kinase kinase 3, MEKKK 3, MAP4K3, RAB8IPL1

Target/Specificity

This MAP4K3 antibody is generated from rabbits immunized with a recombinant protein of human MAP4K3.

Dilution

WB~~1:1000
IHC-P~~1:10~50
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MAP4K3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

MAP4K3 Antibody - Protein Information**Name** MAP4K3**Synonyms** RAB8IPL1

Function May play a role in the response to environmental stress. Appears to act upstream of the JUN N-terminal pathway.

Tissue Location


Ubiquitously expressed in all tissues examined, with high levels in heart, brain, placenta, skeletal muscle, kidney and pancreas and lower levels in lung and liver

MAP4K3 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


MAP4K3 Antibody - Images



Anti-MAP4K3 Antibody at 1:1000 dilution + U-87 MG whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 101 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



MAP4K3 Antibody (Cat. #AP9860a) IHC analysis in formalin fixed and paraffin embedded brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the MAP4K3 Antibody for immunohistochemistry. Clinical relevance has not been evaluated.



MAP4K3 Antibody (Cat. #AP9860a) flow cytometric analysis of HepG2 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

MAP4K3 Antibody - Background

MAP4K3 encodes a member of the Ste20 family of serine/threonine protein kinases. The protein belongs to the subfamily that consists of members, such as germinal center kinase (GCK), that are characterized by an N-terminal catalytic domain and C-terminal regulatory domain. The kinase activity of the encoded protein can be stimulated by UV radiation and tumor necrosis factor- α . The protein specifically activates the c-Jun N-terminal kinase (JNK) signaling pathway. Evidence suggests that it functions upstream of mitogen-activated protein kinase kinase kinase 1 (MEKK1). This gene previously was referred to as RAB8-interacting protein-like 1 (RAB8IPL1), but it has been renamed mitogen-activated protein kinase kinase kinase 3 (MAP4K3).

MAP4K3 Antibody - References

Yan, L., et al. Mol. Cell 37(5):633-642(2010)
Ichikawa, S., et al. J. Bone Miner. Res. (2010) In press :
Lam, D., et al. Proc. Natl. Acad. Sci. U.S.A. 106(29):11978-11983(2009)
Wu, C., et al. Proteomics 7(11):1775-1785(2007)