

**CD160 Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP9869c****Specification**

---

**CD160 Antibody (Center) - Product Information**

Application	WB,E
Primary Accession	<a href="#">O95971</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	19810
Antigen Region	56-82

**CD160 Antibody (Center) - Additional Information****Gene ID** 11126**Other Names**

CD160 antigen, Natural killer cell receptor BY55, CD160, CD160, BY55

**Target/Specificity**

This CD160 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 56-82 amino acids from the Central region of human CD160.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CD160 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**CD160 Antibody (Center) - Protein Information****Name** CD160 {ECO:0000303|PubMed:16809620, ECO:0000312|HGNC:HGNC:17013}

**Function** [CD160 antigen]: Receptor on immune cells capable to deliver stimulatory or inhibitory signals that regulate cell activation and differentiation. Exists as a GPI-anchored and as a transmembrane form, each likely initiating distinct signaling pathways via phosphoinositol

3-kinase in activated NK cells and via LCK and CD247/CD3 zeta chain in activated T cells (PubMed:[19109136](#), PubMed:[11978774](#), PubMed:[17307798](#)). Receptor for both classical and non-classical MHC class I molecules (PubMed:[9973372](#), PubMed:[12486241](#)). In the context of acute viral infection, recognizes HLA-C and triggers NK cell cytotoxic activity, likely playing a role in anti-viral innate immune response (PubMed:[12486241](#)). On CD8+ T cells, binds HLA-A2-B2M in complex with a viral peptide and provides a costimulatory signal to activated/memory T cells (PubMed:[9973372](#)). Upon persistent antigen stimulation, such as occurs during chronic viral infection, may progressively inhibit TCR signaling in memory CD8+ T cells, contributing to T cell exhaustion (PubMed:[25255144](#)). On endothelial cells, recognizes HLA-G and controls angiogenesis in immune privileged sites (PubMed:[16809620](#)). Receptor or ligand for TNF superfamily member TNFRSF14, participating in bidirectional cell-cell contact signaling between antigen presenting cells and lymphocytes. Upon ligation of TNFRSF14, provides stimulatory signal to NK cells enhancing IFNG production and anti-tumor immune response (By similarity). On activated CD4+ T cells, interacts with TNFRSF14 and down-regulates CD28 costimulatory signaling, restricting memory and alloantigen-specific immune response (PubMed:[18193050](#)). In the context of bacterial infection, acts as a ligand for TNFRSF14 on epithelial cells, triggering the production of antimicrobial proteins and pro-inflammatory cytokines (By similarity).

### Cellular Location

[CD160 antigen]: Cell membrane; Lipid-anchor, GPI-anchor

### Tissue Location

Expression is restricted to functional NK and cytotoxic T lymphocytes. Expressed in viral-specific effector memory and terminally differentiated effector memory CD8+ T cells. Expressed in memory and activated CD4+ T cell subsets (at protein level) (PubMed:9743336, PubMed:18193050, PubMed:11978774, PubMed:25255144) Expressed at high levels in intraepithelial lymphocytes (at protein level) (PubMed:9743336). Expressed in both alpha-beta and gamma-delta CD8+ T cell subsets (at protein level) (PubMed:9743336, PubMed:18193050, PubMed:11978774). Expressed in umbilical vein endothelial cells (at protein level) (PubMed:16809620). Expressed in monocytes and at lower levels in B cells (PubMed:23761635). Isoform 3: Expressed exclusively in activated NK cells (at protein level) (PubMed:19109136).


## CD160 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## CD160 Antibody (Center) - Images





Western blot analysis of CD160 Antibody (Center) (Cat. #AP9869c) in Jurkat cell line lysates (35ug/lane). CD160 (arrow) was detected using the purified Pab.

### CD160 Antibody (Center) - Background

CD160 is an 27 kDa glycoprotein which was initially identified with the monoclonal antibody BY55. Its expression is tightly associated with peripheral blood NK cells and CD8 T lymphocytes with cytolytic effector activity. The cDNA sequence of CD160 predicts a cysteine-rich, glycosylphosphatidylinositol-anchored protein of 181 amino acids with a single Ig-like domain weakly homologous to KIR2DL4 molecule. CD160 is expressed at the cell surface as a tightly disulfide-linked multimer. RNA blot analysis revealed CD160 mRNAs of 1.5 and 1.6 kb whose expression was highly restricted to circulating NK and T cells, spleen and small intestine. Within NK cells CD160 is expressed by CD56dimCD16+ cells whereas among circulating T cells its expression is mainly restricted to TCRgd bearing cells and to TCRab+CD8brightCD95+CD56+CD28-CD27-cells. In tissues, CD160 is expressed on all intestinal intraepithelial lymphocytes. CD160 shows a broad specificity for binding to both classical and nonclassical MHC class I molecules.

### CD160 Antibody (Center) - References

- Schmitt, C., et al. Genes Immun. 10(7):616-623(2009)
- Kolz, M., et al. PLoS Genet. 5 (6), E1000504 (2009)
- Giustiniani, J., et al. J. Immunol. 182(1):63-71(2009)
- Cai, G., et al. Nat. Immunol. 9(2):176-185(2008)