

DPAGT1 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP9967a

Specification

DPAGT1 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q9H3H5
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	286-315

DPAGT1 Antibody (Center) - Additional Information

Gene ID 1798

Other Names

UDP-N-acetylglucosamine--dolichyl-phosphate N-acetylglucosaminephosphotransferase, GlcNAc-1-P transferase, G1PT, GPT, N-acetylglucosamine-1-phosphate transferase, DPAGT1, DPAGT2

Target/Specificity

This DPAGT1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 286-315 amino acids from the Central region of human DPAGT1.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DPAGT1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

DPAGT1 Antibody (Center) - Protein Information

Name GPT

Function Catalyzes the initial step of dolichol-linked oligosaccharide biosynthesis in N-linked protein glycosylation pathway: transfers GlcNAc-1-P from UDP-GlcNAc onto the carrier lipid dolichyl

phosphate (P-dolichol), yielding GlcNAc-P-P-dolichol.

Cellular Location


Endoplasmic reticulum membrane; Multi-pass membrane protein

DPAGT1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DPAGT1 Antibody (Center) - Images



Western blot analysis of DPAGT1 Antibody (Center) (Cat. #AP9967a) in MDA-MB231 cell line lysates (35ug/lane). DPAGT1 (arrow) was detected using the purified Pab.

DPAGT1 Antibody (Center) - Background

DPAGT1 encoded by this gene is an enzyme that catalyzes the first step in the dolichol-linked oligosaccharide pathway for glycoprotein biosynthesis. This enzyme belongs to the glycosyltransferase family 4. This protein is an integral membrane protein of the endoplasmic reticulum. The congenital disorder of glycosylation type Ij is caused by mutation in the gene encoding this enzyme.

DPAGT1 Antibody (Center) - References

Nita-Lazar, M., et al. Cancer Res. 69(14):5673-5680(2009)
Bretthauer, R.K. Curr Drug Targets 10(6):477-482(2009)
Wu, X., et al. Hum. Mutat. 22(2):144-150(2003)

DPAGT1 Antibody (Center) - Citations

- [Congenital Myasthenic Syndrome caused by mutations in DPAGT.](#)