

DRAK2 Antibody
Catalog # ASC10073**Specification**

DRAK2 Antibody - Product Information

Application	WB, ICC
Primary Accession	O94768
Other Accession	AB011421 , 3834355
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	DRAK2 antibody can be used for detection of DRAK2 by Western blot 0.5 µg/mL. An approximately 45 kDa band can be detected. Antibody can also be used for immunocytochemistry starting at 10 µg/mL.

DRAK2 Antibody - Additional InformationGene ID **9262****Other Names**

DRAK2 Antibody: DRAK2, DRAK2, DAP kinase-related apoptosis-inducing protein kinase 2, serine/threonine kinase 17b

Target/Specificity

STK17B; It has no cross responses to DAP or ZIP kinases. The approximately 70 kDa band is probably non-related to DRAK2 although it is peptide blockable.

Reconstitution & Storage

DRAK2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

DRAK2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

DRAK2 Antibody - Protein Information**Name** STK17B**Synonyms** DRAK2**Function**

Phosphorylates myosin light chains (By similarity). Acts as a positive regulator of apoptosis.

Cellular Location

Nucleus. Cell membrane. Endoplasmic reticulum-Golgi intermediate compartment.
Note=Colocalizes with STK17B at the plasma membrane.


Tissue Location

Highly expressed in placenta, lung, pancreas. Lower levels in heart, brain, liver, skeletal muscle and kidney


DRAK2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DRAK2 Antibody - Images

Western blot analysis of DRAK2 in Raji cell lysate with DRAK2 antibody at (A) 1 and B (2) μ g/mL.



Immunocytochemistry of DRAK2 in Jurkat cells with DRAK2 antibody at 10 μ g/mL.

DRAK2 Antibody - Background

DRAK2 Antibody: Apoptosis is mediated by death domain containing adapter molecules and a

caspase family of proteases. Certain serine/threonine protein kinases, such as ASK-1 and RIP, are mediators of apoptosis. Two novel serine/threonine kinases that induce apoptosis were recently identified and designated DRAK1 and DRAK2 (for DAP kinase-related apoptosis-inducing protein kinases). DRAKs contain an N-terminal kinase domain and a C-terminal regulation domain. Overexpression of DRAK2 induces apoptosis. DRAKs have high sequence homology to DAP and ZIP kinases, and they represent a novel family of serine/threonine kinases, which mediates apoptosis through their catalytic activities. DRAK2 is located in nucleus and the messenger RNA was ubiquitously expressed in human tissues.

DRAK2 Antibody - References

Sanjo H, Kawai T, Akira S. DRAKs, novel serine/threonine kinases related to death-associated protein kinase that trigger apoptosis. J Biol Chem 1998;273:29066-71 (RD1299)