

DFF40 Antibody
Catalog # ASC10074**Specification**

DFF40 Antibody - Product Information

Application	WB, ICC, IF
Primary Accession	O54788
Other Accession	NP_031885 , 160948620
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	40 kDa KDa
Application Notes	DFF40 antibody can be used for detection of DFF40 by Western blot at 0.5 µg/mL. A 40 kDa band can be detected. Antibody can also be used for immunocytochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL.

DFF40 Antibody - Additional InformationGene ID **13368****Other Names**

DFF40 Antibody: CAD, CPAN, 40kDa, DFF40, Didff, 5730477D02Rik, Cad, DNA fragmentation factor subunit beta, Caspase-activated deoxyribonuclease, CAD, DNA fragmentation factor, beta subunit

Target/Specificity

Dfffb;

Reconstitution & Storage

DFF40 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

DFF40 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

DFF40 Antibody - Protein Information**Name** Dfffb**Synonyms** Cad**Function**

Nuclease that induces DNA fragmentation and chromatin condensation during apoptosis. Degrades naked DNA and induces apoptotic morphology.


Cellular Location

Cytoplasm. Nucleus.


DFF40 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DFF40 Antibody - Images

Western blot analysis of DFF40 in Jurkat (J) and K562 (K) whole cell lysate with DFF40 antibody at 1:500 dilution.



Immunocytochemistry of DFF40 in Jurkat cells with DFF40 antibody at 5 µg/mL.



Immunofluorescence of DFF40 in Jurkat cells with DFF40 antibody at 20 µg/mL.

DFF40 Antibody - Background

DFF40 Antibody: Apoptosis is related to many diseases and induced by a family of cell death receptors and their ligands. Cell death signals are transduced by death domain containing adapter molecules and members of the caspase family of proteases. These death signals finally cause the degradation of chromosomal DNA by activated DNase. A mouse DNase that causes DNA fragmentation was identified recently and designated CAD for caspase activated deoxyribonuclease. The human homologue of mouse CAD was more recently identified by three groups independently and termed CPAN, DFF40, and human CAD, respectively. DFF45/ICAD is the inhibitory protein of DFF40/CAD and forms complex with DFF40/CAD. Upon cleavage of DFF45/ICAD by activated caspase, DFF40/CAD is released and activated and eventually causes the degradation of DNA in the nuclei. Activation of DFF40/CAD, which causes DNA degradation, is the hallmark of apoptotic cell death.

DFF40 Antibody - References

Enari M, Sakahira H, Yokoyama H, Okawa K, Iwamatsu A, Nagata S. A caspase-activated DNase that degrades DNA during apoptosis, and its inhibitor ICAD. *Nature* 1998;391:43-50
Sakahira H, Enari M, Nagata S. Cleavage of CAD inhibitor in CAD activation and DNA degradation during apoptosis. *Nature* 1998;391:96-99
Halenbeck R, MacDonald H, Roulston A, Chen TT, Conroy L, Williams LT. CPAN, a human nuclease regulated by the caspase-sensitive inhibitor DFF45. *Curr Biol* 1998;8:537-40
Liu X, Li P, Widlak P, Zou H, Luo X, Garrard WT, Wang X The 40-kDa subunit of DNA fragmentation factor induces DNA fragmentation and chromatin condensation during apoptosis. *Proc Natl Acad Sci USA* 1998;95:8461-6