

**CSN8 Antibody**  
**Catalog # ASC10691****Specification**

---

**CSN8 Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q99627</a>
Other Accession	<a href="#">NP_006701</a> , <a href="#">5729779</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	CSN8 antibody can be used for detection of CSN8 by Western blot at 2 µg/mL.

**CSN8 Antibody - Additional Information**

Gene ID	10920
Target/Specificity	
COPS8;	

**Reconstitution & Storage**

CSN8 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

**Precautions**

CSN8 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**CSN8 Antibody - Protein Information**

**Name** COPS8

**Synonyms** CSN8

**Function**

Component of the COP9 signalosome complex (CSN), a complex involved in various cellular and developmental processes. The CSN complex is an essential regulator of the ubiquitin (Ubl) conjugation pathway by mediating the deneddylation of the cullin subunits of SCF- type E3 ligase complexes, leading to decrease the Ubl ligase activity of SCF-type complexes such as SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, c-jun/JUN, IκappaBα/NFKBIA, ITPK1 and IRF8/ICSBP, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the Ubl system, respectively.

**Cellular Location**


Cytoplasm. Nucleus

## CSN8 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## CSN8 Antibody - Images



Western blot analysis of CSN8 in Human liver lysate with CSN8 antibody at 2 µg/mL.

## CSN8 Antibody - Background

**CSN8 Antibody:** The COP9 signalosome (CSN) is an evolutionarily conserved protein complex of the eight subunits that interacts with deubiquitinating enzymes and protein kinases and is highly homologous to the lid sub-complex of 26S proteasome. The CSN complex is an essential regulator of the ubiquitin conjugation pathway by mediating the deneddylation of the SCF-type E3 ligase complexes, which leads to a decrease in ubiquitin ligase activity of SCF-complexes such as SCF, CSA or DDB2. It is also involved in phosphorylation of p53, c-jun/JUN, ITPK1 and IRF8/ICSBP, possibly via its association with CK2 and PKD kinases. CSN8 encodes the smallest and the least conserved but first identified subunit of CSN. Recent studies show CSN8 is essential for *Drosophila* development and is essential for peripheral T cell homeostasis and antigen receptor-induced entry into the cell cycle from quiescence.

## CSN8 Antibody - References

Wei N and Deng XW. The COP9 signalosome. *Annu. Rev. Cell Dev. Biol.*2003; 19:261-86.  
Bech-Otschir D, Seeger M, and Dubiel W. The COP9 signalosome: at the interface between signal transduction and ubiquitin-dependent proteolysis. *J. Cell Sci.*2002; 115:467-73.  
Groisman R, Polanowska J, Kuraoka I, et al. The ubiquitin ligase activity in the DDB2 and regulated by the COP9 signalosome in response to DNA damage. *Cell*2003; 113:357-67.

Uhle S, Medalia O, Waldron R, et al. Protein kinase CK2 and protein kinase D are associated with the COP9 signalosome. EMBO J.2003; 22:1302-12.