

# **ZCRB1** Antibody

Catalog # ASC10926

### **Specification**

# **ZCRB1 Antibody - Product Information**

Application
Primary Accession
Other Accession
Reactivity
Host
Clonality

Clonality Isotype Application Notes WB

<u>Q8TBF4</u> <u>NP\_149105</u>, <u>51243065</u>

Human, Mouse, Rat Rabbit Polyclonal

IgG

ZCRB1 antibody can be used for detection of ZCRB1 by Western blot at 1 - 2 μg/mL.

# **ZCRB1** Antibody - Additional Information

Gene ID **85437** 

Target/Specificity

ZCRB1;

### **Reconstitution & Storage**

ZCRB1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

#### **Precautions**

ZCRB1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

### **ZCRB1 Antibody - Protein Information**

Name ZCRB1

**Cellular Location** Nucleus, nucleoplasm

# **ZCRB1 Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation



- Flow Cytomety
- Cell Culture

# **ZCRB1 Antibody - Images**



Western blot analysis of ZCRB1 in Raji cell lysate with ZCRB1 antibody at (A) 1 and (B) 2 µg/mL.

# **ZCRB1 Antibody - Background**

ZCRB1 Antibody: ZCRB1 is a nuclear protein first identified in a differential display screen involving morphine-dependence-related genes. Its expression is increased following morphine treatment and has been observed to be elevated in HepG2 cells. ZCRB1 contains a CCHC-type zing finger RNA-binding motif and can interact with the DBA-binding domain of the stem cell regulator C/EBP transcription factors. ZCRB1 was identified as one of the protein components of U11/U12 snRNPs, which are components of U12-type spliceosome and function as a molecular bridge connecting both ends of the intron, suggesting ZCRB1 may play a key role in U12-type splicing.

### **ZCRB1 Antibody - References**

Wang H, Gao MX, Li L, et al. Isolation, expression, and characterization of the human ZCRB1 gene mapped to 12g2. Genomics2007; 89:59-69.

Trivedi AK, Bararia D, Christopeit M, et al. Proteomic identification of C/EBP-DBD multiprotein complex: JNK1 activates stem cell regulator C/EBPalpha by inhibiting its ubiquitination. Oncogene2007; 1789-801.

Will CL, Schneider C, Hossbach M, et al. The human 18S U11/U12 snRNP contains a set of novel proteins not found in the U2-dependent spliceosome. RNA2004; 10:929-41.