

# THEM2 Antibody

Catalog # ASC10940

#### Specification

## THEM2 Antibody - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Application Notes WB, IHC, IF <u>O9NPJ3</u> NP\_060943, 8923812 Human, Mouse, Rat Rabbit Polyclonal IgG THEM2 antibody can be used for detection of THEM2 by Western blot at 1 μg/mL. Antibody can also be used for immunohistochemistry starting at 2.5 μg/mL. For immunofluorescence start at 20 μg/mL.

### THEM2 Antibody - Additional Information

Gene ID Target/Specificity ACOT13;

#### **Reconstitution & Storage**

THEM2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

55856

#### Precautions

THEM2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

### **THEM2 Antibody - Protein Information**

Name ACOT13 (HGNC:20999)

#### Synonyms THEM2

Function

Catalyzes the hydrolysis of acyl-CoAs into free fatty acids and coenzyme A (CoASH), regulating their respective intracellular levels (PubMed:<a href="http://www.uniprot.org/citations/16934754" target="\_blank">16934754</a>, PubMed:<a href="http://www.uniprot.org/citations/19170545" target="\_blank">16934754</a>, PubMed:<a href="http://www.uniprot.org/citations/19170545" target="\_blank">19170545</a>). Has acyl-CoA thioesterase activity towards medium (C12) and long-chain (C18) fatty acyl-CoA substrates (By similarity) (PubMed:<a

href="http://www.uniprot.org/citations/16934754" target="\_blank">16934754</a>, PubMed:<a href="http://www.uniprot.org/citations/19170545" target="\_blank">19170545</a>). Can also hydrolyze 3-hydroxyphenylacetyl-CoA and 3,4-dihydroxyphenylacetyl-CoA (in vitro) (By similarity)



(PubMed:<a href="http://www.uniprot.org/citations/16934754" target="\_blank">16934754</a>, PubMed:<a href="http://www.uniprot.org/citations/19170545" target="\_blank">19170545</a>). May play a role in controlling adaptive thermogenesis (By similarity).


#### **Cellular Location**

Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q9CQR4}. Mitochondrion {ECO:0000250|UniProtKB:Q9CQR4}. Nucleus {ECO:0000250|UniProtKB:Q9CQR4} Cytoplasm, cytoskeleton, spindle {ECO:0000250|UniProtKB:Q9CQR4} Note=During interphase, found both in the nucleus and in the cytoplasm At mitosis, localizes to the spindle. Colocalizes with tubulin {ECO:0000250|UniProtKB:Q9CQR4}

#### **THEM2 Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>
- THEM2 Antibody Images




Western blot analysis of THEM2 in HepG2 cell lysate with THEM2 antibody at 1  $\mu$ g/mL in (A) the absence and (B) the presence of blocking peptide.





Immunohistochemistry of THEM2 in human liver tissue with THEM2 antibody at 2.5 µg/mL.



Immunofluorescence of THEM2 in human liver tissue with THEM2 antibody at 20 µg/mL.

### THEM2 Antibody - Background

THEM2 Antibody: THEM2, also known as ACOT13, belongs to the hotdog-fold superfamily and possesses thioesterase activity, with greater activity observed with longer chain acyl-CoAs such as myristoyl- and palmitoyl-CoA. THEM2 is highly expressed in several tissues such as heart, kidney, liver and brain and has been shown to be co-localized with beta-tubulin on microtubules. THEM2 interacts with StarD, a protein that plays a key role in fatty acid metabolism, and the addition of StarD to THEM2 increased its catalytic activity, suggesting that THEM2 plays a significant role in the metabolism of fatty acids. At least two isoforms of THEM2 are known to exist.

### **THEM2 Antibody - References**

Cheng Z, Song F, Shan X, et al. Crystal structure of human thioesterase superfamily member 2. Biochem. Biophys. Res. Commun.2006; 349:172-7.

Wei J, Kang HW, and Cohen DE. Thioesterase superfamily member 2 (Them2)/acyl-CoA thioesterase 13 (Acot13): a homotetrameric hotdog fold thioesterase with selectivity for long-chain fatty acyl-CoAs. Biochem. J.2009; 421:311-22.

Cheng Z, Bao S, Shan X, et al. Human thioesterase superfamily member 2 (hTHEM2) is co-localized with beta-tubulin onto the microtubule. Biochem. Biophys. Res. Commun.2006; 350:850-3. Scappa EF, Pocai A, Wu MK, et al. Regulation of energy substrate utilization and hepatic insulin sensitivity by phosphatidylcholine transfer protein/StarD. FASEB J.2008; 22:2579-90.