

## LYRM3 Antibody

Catalog # ASC11093

## Specification

## **LYRM3 Antibody - Product Information**

Application
Primary Accession
Other Accession
Reactivity
Host
Clonality
Isotype

Isotype Application Notes WB, IHC, IF <u>Q9Y6M9</u>

<u>09Y6M9</u>, <u>8134589</u> **Human, Mouse, Rat** 

Rabbit Polyclonal

IgG

LYRM3 antibody can be used for detection of LYRM3 by Western blot at 1 - 2  $\mu$ g/mL.

Antibody can also be used for

immunohistochemistry starting at 5  $\mu$ g/mL. For immunofluorescence start at 20  $\mu$ g/mL.

# LYRM3 Antibody - Additional Information

Gene ID 4715

**Target/Specificity** 

NDUFB9;

### **Reconstitution & Storage**

LYRM3 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

#### **Precautions**

LYRM3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

#### LYRM3 Antibody - Protein Information

#### Name NDUFB9

Synonyms LYRM3, UQOR22

### **Function**

Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.

# **Cellular Location**

Mitochondrion inner membrane; Peripheral membrane protein; Matrix side




# **LYRM3 Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# **LYRM3 Antibody - Images**




Western blot analysis of LYRM3 in human liver tissue lysate with LYRM3 antibody at (A) 1 and (B) 2  $\mu g/mL$ .



Immunohistochemistry of LYRM3 in rat liver tissue with LYRM3 antibody at 5 μg/mL.





Immunofluorescence of LYRM3 in rat liver tissue with LYRM3 antibody at 20 µg/mL.

## LYRM3 Antibody - Background

LYRM3 Antibody: LYRM3, also known as NADH dehydrogenase (ubiquinone) 1 beta subcomplex 9 (NDUFB9), is a ubiquitously expressed LYR-motif containing protein. It has been suggested to be a candidate gene for the branchio-oto-renal (BOR) syndrome, which is characterized by branchial and renal abnormalities and heredity deafness disorders. Other than its LYR-motif, LYRM3 appears to have no functional or structural relationship to either LYRM1 or LYRM2.

## **LYRM3 Antibody - References**

Lin X, Wells DE, Kimberling WJ, et al. Human NDUFB9 gene: genomic organization and a possible candidate gene associated with deafness disorder mapped to chromosome 8q13. Hum. Heredity1999; 49:75-80.