

TMEM184C Antibody

Catalog # ASC11104

Specification

TMEM184C Antibody - Product Information

Application
Primary Accession
Other Accession
Reactivity
Host
Clonality

Isotype

Application Notes

WB, IHC, IF O9NVA4

NP_060711, 190358512 Human, Mouse, Rat

Rabbit Polyclonal

lgG

TMEM184C antibody can be used for detection of TMEM184C by Western blot at 1 μg/mL. Antibody can also be used for immunohistochemistry starting at 5 μg/mL. For immunofluorescence start at 20 μg/mL.

TMEM184C Antibody - Additional Information

Gene ID **55751**

Target/Specificity

TMEM184C;

Reconstitution & Storage

TMEM184C antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

TMEM184C Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

TMEM184C Antibody - Protein Information

Name TMEM184C

Synonyms TMEM34

Function

Possible tumor suppressor which may play a role in cell growth.

Cellular Location

Membrane; Multi-pass membrane protein

Tissue Location

Widely expressed with higher expression in lung, kidney, spleen, pancreas, thymus, prostate, testis, ovary, small intestine and thyroid.




TMEM184C Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

TMEM184C Antibody - Images




Western blot analysis of TMEM184C in mouse heart tissue lysate with TMEM184C antibody at 1 $\mu g/mL$.



Immunohistochemistry of TMEM184C in mouse heart tissue with TMEM184C antibody at 5 $\mu g/mL$.





Immunofluorescence of TMEM184C in mouse heart tissue with TMEM184B antibody at 20 $\mu g/mL$.

TMEM184C Antibody - Background

TMEM184C Antibody: Anaplastic thyroid cancer is one of the most lethal forms of cancer, but the precise carcinogenic mechanism has not been identified. TMEM184C, also known as TMEM34, was identified in a cDNA microarray analysis as being down-regulated in anaplastic thyroid cancers compared to normal thyroid tissues. TMEM184C protein expression was also lower in cell lines derived from these types of cancers compared to that of normal thyroid tissues or cell lines based on other types of thyroid cancers. Furthermore, transfection of TMEM34 into KTA2 cells led to the inhibition of cell growth, suggesting that TMEM184C might act as a tumor suppressor in anaplastic thyroid cancers.

TMEM184C Antibody - References

Akaishi J, Onda M, Okamoto J, et al. Down-regulation of an inhibitor of cell growth, transmembrane protein 34 (TMEM34), in anaplastic thyroid cancer. J. Cancer Res. Clin. Oncol.2007; 133:213-8.