10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

APC8 Antibody
Catalog # ASC11120

Specification

APC8 Antibody - Product Information

Application	**WB, ICC**
Primary Accession	Q9UJX2
Other Accession	Q9UJX2, 254763423
Reactivity	**Human, Mouse, Rat**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**IgG**
Application Notes	**APC8 antibody can be used for detection of APC8 by Western blot at 1 - 2 µg/mL. Antibody can also be used for immunocytochemistry starting at 5 µg/mL.**

APC8 Antibody - Additional Information

Gene ID	**8697**

Target/Specificity
CDC23;

Reconstitution & Storage
APC8 antibody can be stored at 4℃ for three months and -20℃, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions
APC8 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

APC8 Antibody - Protein Information

Name CDC23

Synonyms ANAPC8

Function
Component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle. The APC/C complex acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains.


APC8 Antibody - Protocols

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

APC8 Antibody - Images



Western blot analysis of APC8 in K562 cell lysate with APC8 antibody at (A) 1 and (B) 2 µg/mL.



Immunocytochemistry of APC8 in K562 cells with APC8 antibody at 5 µg/mL.

APC8 Antibody - Background

APC8 Antibody: Cell cycle regulated protein ubiquitination and degradation within subcellular domains is thought to be essential for the normal progression of mitosis. APC8 is a highly conserved component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle. APC/C is responsible for degrading anaphase inhibitors, mitotic cyclins, and spindle-associated proteins ensuring that events of mitosis take place in proper sequence. The individual APC/C components mRNA and protein levels are expressed at approximately the same levels in most tissues and cell

lines, suggesting that they perform their functions as part of a complex. In Drosophila, silencing of APC8 results in developmental delay and pupal lethality with elevated levels of apoptosis, high mitotic index, and delayed or blocked mitosis.

APC8 Antibody - References

JM Peters. The anaphase promoting complex/cyclosome: a machine designed to destroy. Nat. Rev. Mol. Cell Biol.2006; 7:644-56.
Jorgensen PM, Graslund S, Betz R, et al. Characterisation of the human APC1, the largest subunit of the anaphase-promoting complex. Gene2001; 262:51-9.
Pal M, Nagy O, Menesi D, et al. Structurally related TPR subunits contribute differently to the function of the anaphase-promoting complex in Drosophila melanogaster. J. Cell Sci.2007; 120:3238-48.