

TCTN2 Antibody
Catalog # ASC11162**Specification**

TCTN2 Antibody - Product Information

Application	WB
Primary Accession	Q96GX1
Other Accession	NP_079085 , 31377681
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	TCTN2 antibody can be used for detection of TCTN2 by Western blot at 1 µg/mL.

TCTN2 Antibody - Additional Information

Gene ID	79867
Target/Specificity	
TCTN2;	

Reconstitution & Storage

TCTN2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

TCTN2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

TCTN2 Antibody - Protein Information

Name TCTN2

Synonyms C12orf38, TECT2

Function

Component of the tectonic-like complex, a complex localized at the transition zone of primary cilia and acting as a barrier that prevents diffusion of transmembrane proteins between the cilia and plasma membranes. Required for hedgehog signaling transduction (By similarity).

Cellular Location


Membrane; Single-pass type I membrane protein. Cytoplasm, cytoskeleton, cilium basal body. Note=Localizes at the transition zone, a region between the basal body and the ciliary axoneme.

TCTN2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TCTN2 Antibody - Images



Western blot analysis of TCTN2 in SK-N-SH cell lysate with TCTN2 antibody at 1 μ g/mL.

TCTN2 Antibody - Background

TCTN2 Antibody: TCTN2 is a member of the Tectonic protein family, a group of evolutionarily conserved secreted and transmembrane proteins that regulate the Hedgehog (Hh)-mediated patterning of the neural tube. While the expression pattern and function of TCTN2 is not known, it is 49% similar to TCTN1. TCTN1 is expressed during embryonic development in regions that participate in Hh signaling, beginning in the gastrulation stages in the ventral node. Mice expressing mutant TCTN1 die between E13.5 and E16.5 and display holoprosencephaly, a defect associated with reduced Hh signaling, indicating the role of TCTN1 as an Hh activator. At later stages in development, TCTN1 is thought to also act as a repressor on the Hh pathway in the anterior and posterior neural tube.

TCTN2 Antibody - References

Reiter JF and Skarnes WC. Tectonic, a novel regulator of the Hedgehog pathway required for both activation and inhibition. *Genes Dev.*2006; 20:22-7.