

KPNA1 Antibody

Catalog # ASC11205

Specification

KPNA1 Antibody - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Application Notes WB, ICC, IF <u>P52294</u> NP_002255, 222144293 Human, Mouse, Rat Rabbit Polyclonal IgG KPNA1 antibody can be used for detection of KPNA1 by Western blot at 1 μg/mL. Antibody can also be used for immunocytochemistry starting at 2.5 μg/mL. For immunofluorescence start at 20 μg/mL.

KPNA1 Antibody - Additional Information

Gene ID Target/Specificity KPNA1;

Reconstitution & Storage

KPNA1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

3836

Precautions

KPNA1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

KPNA1 Antibody - Protein Information

Name KPNA1

Synonyms RCH2

Function

Functions in nuclear protein import as an adapter protein for nuclear receptor KPNB1 (PubMed:7892216, PubMed:8692858, PubMed:27713473). Binds specifically and directly to substrates containing either a simple or bipartite NLS motif (PubMed:7892216, PubMed:7892216, PubMed:7892216, PubMed:8692858, PubMed:27713473, PubMed:27713473, PubMed:27713473, PubMed:27713473, PubMed:27713473, DubMed:27713473, DubMed:<



the importin/substrate complex to the nuclear pore complex (NPC) is mediated by KPNB1 through binding to nucleoporin FxFG repeats and the complex is subsequently translocated through the pore by an energy requiring, Ran-dependent mechanism (PubMed:7892216, PubMed:27713473). At the nucleoplasmic side of the NPC, Ran binds to importin-beta and the three components separate and importin-alpha and -beta are re-exported from the nucleus to the cytoplasm where GTP hydrolysis releases Ran from importin (PubMed:7892216). The directionality of nuclear import is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus (PubMed:7892216).

Cellular Location Cytoplasm. Nucleus


Tissue Location Expressed ubiquitously.

KPNA1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

KPNA1 Antibody - Images




Western blot analysis of KPNA1 in Hela cell lysate with KPNA1 antibody at 1 μ g/mL.





Immunocytochemistry of KPNA1 in HeLa cells with KPNA1 antibody at 2.5 µg/mL.



Immunofluorescence of KPNA1 in K562 cells with KPNA1 antibody at 20 $\mu\text{g/mL}.$

KPNA1 Antibody - Background

KPNA1 Antibody: Karyopherin, a cytosolic and heterodimeric protein complex consisting of alpha and beta subunits, is responsible for targeting proteins with nuclear localization signals to the nuclear pore complex by an energy requiring, Ran-dependent mechanism. The alpha subunit and imported substrate enter the nucleus and accumulate in the nucleoplasm, while the beta subunit accumulates at the NPC. KPNA1, also known as importin alpha 5, is the alpha subunit of karyopherin, which forms a complex with importin subunit beta-1 and functions in nuclear protein import as an adapter protein for nuclear receptor KPNB1. It is ubiquitously expressed and polyubiquitinated in the presence of RAG1. KPNA1 interacts with various virus nucleoproteins, including those of Ebola and influenza.

KPNA1 Antibody - References

Moroianu J. Molecular mechanisms of nuclear protein transport. Crit. Rev. Eukaryot. Gene Expr.1997; 7:61-72.

Gilchrist D and Rexach M. Molecular basis for the rapid dissociation of nuclear localization signals from karyopherin alpha in the nucleoplasm. J. Biol. Chem.2003; 278: 51937-49.

Simkus C, Makiya M and Jones JM. Karyopherin alpha 1 is a putative substrate of the RAG1 ubiquitin ligase. Mol. Immunol.2009; 46:1319-25.

Reid SP, Valmas C, and Martinez O. Ebola virus VP24 proteins inhibit the interaction of NPI-1 subfamily karyopherin alpha proteins with activated STAT1. J. Virol.2007; 81:13469-77.