

Anosmin Antibody
Catalog # ASC11330**Specification**

Anosmin Antibody - Product Information

Application	WB, ICC, IF
Primary Accession	P23352
Other Accession	NP_000207 , 119395746
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	Anosmin antibody can be used for detection of Anosmin by Western blot at 1 µg/mL. Antibody can also be used for immunocytochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL.

Anosmin Antibody - Additional Information

Gene ID	3730
Target/Specificity	
KAL1;	

Reconstitution & Storage

Anosmin antibody can be stored at 4 °C, stable for one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

Anosmin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anosmin Antibody - Protein Information

Name ANOS1 ([HGNC:6211](#))

Synonyms ADMLX, KAL, KAL1, KALIG1

Function

Has a dual branch-promoting and guidance activity, which may play an important role in the patterning of mitral and tufted cell collaterals to the olfactory cortex (By similarity).
Chemoattractant for fetal olfactory epithelial cells.

Cellular Location

Cell membrane; Peripheral membrane protein. Secreted. Note=Proteolytic cleavage may release it from the cell surface into the extracellular space

Tissue Location


Expressed in the cerebellum (at protein level).

Anosmin Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anosmin Antibody - Images



Western blot analysis of Anosmin in MCF7 cell lysate with Anosmin antibody at 1 μ g/mL.



Immunocytochemistry of Anosmin in MCF7 cells with Anosmin antibody at 5 μ g/mL.



Immunofluorescence of Anosmin in MCF7 cells with Anosmin antibody at 20 µg/mL.

Anosmin Antibody - Background

Anosmin Antibody: Mutations in Anosmin-1, an extracellular matrix-associated glycosylated protein, have been linked with Kallmann Syndrome (KS), an X-linked genetic disorder characterized by loss of smell caused by abnormal olfactory bulb development and delayed puberty caused by disrupted migration of the gonadotropin-releasing hormone neuron. Anosmin-1 has been shown to directly bind FGFR1 via its N-terminal cysteine-rich domain, whey-acidic protein-like domain, and its first FnIII repeat with the D2 and D3 ectodomains of FGFR1. It is thought that Anosmin-1 can modulate FGFR1 signaling and have opposing effects on the formation and activation of FGF2-FGFR1-heparin complex.

Anosmin Antibody - References

Franco B, Guioli S, Pragliola A, et al. A gene deleted in Kallmann's syndrome shares homology with neural cell adhesion and axonal path-finding molecules. *Nature* 1991; 353:529-36.

Soussi-Yanicostas N, Hardelin JP, Arroyo-Jimenez MM, et al. Initial characterization of anosmin-1, a putative extracellular matrix protein synthesized by definite neuronal cell populations in the central nervous system. *J. Cell Sci.* 1996; 109:1749-57.

Hu Y, Guimond SE, Travers P, et al. Novel mechanisms of fibroblast growth factor receptor 1 regulation by extracellular matrix protein Anosmin-1. *J. Biol. Chem.* 2009; 284:29905-20