

UPK1B Antibody

Catalog # ASC11380

Specification

UPK1B Antibody - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Application Notes WB, IHC, IF <u>O75841</u> NP_008883, 49619237 Human, Mouse, Rat Rabbit Polyclonal IgG UPK1B antibody can be used for detection of UPK1B by Western blot at 0.5 - 1 μg/mL. Antibody can also be used for immunohistochemistry starting at 5 μg/mL. For immunofluorescence start at 20 μg/mL.

UPK1B Antibody - Additional Information

Gene ID

7348

Target/Specificity

UPK1B; At least two isoforms of UPK1B are known to exist; this antibody will recognize both isoforms. UPK1B antibody is predicted to not cross-react with other tetraspanin protein family members.

Reconstitution & Storage

UPK1B antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

UPK1B Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

UPK1B Antibody - Protein Information

Name UPK1B

Synonyms TSPAN20

Function

Component of the asymmetric unit membrane (AUM); a highly specialized biomembrane elaborated by terminally differentiated urothelial cells. May play an important role in normal bladder epithelial physiology, possibly in regulating membrane permeability of superficial umbrella cells or in stabilizing the apical membrane through AUM/cytoskeletal interactions (By similarity).

Cellular Location

Membrane; Multi-pass membrane protein.




Tissue Location Bladder epithelium.

UPK1B Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

UPK1B Antibody - Images




Western blot analysis of UPK1B in mouse bladder tissue lysate with UPK1B antibody at 0.5 μ g/mL in the (A) absence and (B) presence of blocking peptide.



Immunohistochemistry of UPK1B in human bladder tissue with UPK1B antibody at 5 µg/mL.





Immunofluorescence of UPK1B in human bladder muscle tissue with UPK1B antibody at 20 µg/mL. **UPK1B Antibody - Background**

UPK1B Antibody: Uroplakin-1B (UPK1B) is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. UPK1B is found in the asymmetrical unit membrane (AUM) where it can form a complex with other transmembrane 4 superfamily proteins and may play a role in normal bladder epithelial physiology, possibly in regulating membrane permeability of superficial umbrella cells or in stabilizing the apical membrane through AUM/cytoskeletal interactions.

UPK1B Antibody - References

Yu J, Lin JH, Wu XR, et al. Uroplakins la and lb, two major differentiation products of bladder epithelium, belong to a family of four transmembrane domain (4TM) proteins. J. Cell Biol. 1994; 125:171-82.

Wu XR, Kong XP, Pellicer A, et al. Uroplakins in urothelial biology, function, and disease. Kidney Int. 2009; 75:1153-65

Sun TT, Zhao H, Provet J, et al. Formation of asymmetric unit membrane during urothelial differentiation. Mol. Biol. Rep. 1996; 23:3-11.