

TSHZ2 Antibody
Catalog # ASC11451**Specification**

TSHZ2 Antibody - Product Information

Application	WB, IF
Primary Accession	Q9NRE2
Other Accession	NP_775756 , 153945734
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	TSHZ2 antibody can be used for detection of TSHZ2 by Western blot at 1 µg/mL. Antibody can also be used for immunofluorescence starting at 20 µg/mL. For immunofluorescence start at 5 µg/mL.

TSHZ2 Antibody - Additional Information

Gene ID	128553
Target/Specificity	
TSHZ2;	

Reconstitution & Storage

TSHZ2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

TSHZ2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

TSHZ2 Antibody - Protein Information

Name TSHZ2

Synonyms C20orf17, TSH2, ZNF218

Function

Probable transcriptional regulator involved in developmental processes. May act as a transcriptional repressor (Potential).

Cellular Location

Nucleus.

Tissue Location


Expressed in brain; strongly reduced in post-mortem elderly subjects with Alzheimer disease.

TSHZ2 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TSHZ2 Antibody - Images



Western blot analysis of TSHZ2 in A-20 cell lysate with TSHZ2 antibody at 1 μ g/mL in (A) the absence and (B) the presence of blocking peptide.



Immunofluorescence of TSHZ2 in A20 cells with TSHZ2 antibody at 20 μ g/mL.

TSHZ2 Antibody - Background

TSHZ2 Antibody: The Teashirt zinc finger homeobox (TSHZ) family comprise a family of evolutionarily conserved transcription factors that, in *Drosophila*, are active in specific body parts for patterning, but whose function in vertebrates is less clear. In mice, the known three TSHZ proteins are expressed in distinct patterns in the developing and adult brain, suggesting that they play a role in the establishment of regional identity and specification of cell types within the brain. Recent experiments have shown that the expression of TSHZ2 is frequently downregulated in most

breast and prostate cancers and its promoter was unmethylated in virtually all cases, suggesting this family of proteins may also be involved in carcinogenesis.

TSHZ2 Antibody - References

Caubit X, Core N, Boned A, et al. Vertebrate orthologues of the Drosophila region-specific patterning gene *teashirt*. *Mech. Dev.* 2000; 91:445-8.

Santos JS, Fonseca NA, Vieira CP, et al. Phylogeny of the Teashirt-related zinc finger (*tshz*) gene family and analysis of the developmental expression of *tshz2* and *tshz3* in the zebrafish. *Dev. Dyn.* 2010; 239:1010-8.

Caubit X, Tiveron MC, Cremer H, et al. Expression patterns of the three Teashirt-related genes define specific boundaries in the developing and postnatal mouse forebrain. *J. Comp. Neurol.* 2005; 486:76-88.

Yamamoto M, Cid E, Bru S, et al. Rare and frequent promoter methylation, respectively, of TSHZ2 and 3 genes that are both downregulated in expression in breast and prostate cancers. *PLoS ONE* 2011; 6:e17149.