10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

LEMD2 Antibody
Catalog # ASC11495

Specification

LEMD2 Antibody - Product Information

Application	**WB, ICC, IF**
Primary Accession	Q8NC56
Other Accession	NP_851853, 31044432
Reactivity	**Human, Mouse**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**IgG**
Application Notes	**LEMD2 antibody can be used for detection of LEMD2 by Western blot at 1 - 2 µg/mL. Antibody can also be used for immunocytochemistry starting at 2.5 µg/mL. For immunofluorescence start at 2.5 µg/mL.**

LEMD2 Antibody - Additional Information

Gene ID **221496**

Target/Specificity
LEMD2; At least two isoforms of LEMD2 are known to exist; this antibody will detect only the longer isoform.

Reconstitution & Storage
LEMD2 antibody can be stored at 4℃ for three months and -20℃, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions
LEMD2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

LEMD2 Antibody - Protein Information

Name LEMD2

Function
Nuclear lamina-associated inner nuclear membrane protein that is involved in nuclear structure organization, maintenance of nuclear envelope (NE) integrity and NE reformation after mitosis (PubMed:16339967, PubMed:17097643, PubMed:28242692, PubMed:32494070). Plays a role as transmembrane adapter for the endosomal sorting complexes required for transport (ESCRT), and is thereby involved in ESCRT-mediated NE reformation (PubMed:28242692, PubMed:<a

href="http://www.uniprot.org/citations/32494070" target="_blank">32494070). Promotes ESCRT-mediated NE closure by recruiting CHMP7 and downstream ESCRT-III proteins IST1/CHMP8 and CHMP2A to the reforming NE during anaphase (PubMed:28242692). During nuclear reassembly, condenses into a liquid-like coating around microtubule spindles and coassembles with CHMP7 to form a macromolecular O-ring seal at the confluence between membranes, chromatin, and the spindle to facilitate early nuclear sealing (PubMed:32494070). Plays a role in the organization of heterochromatin associated with the NE and in the maintenance of NE organization under mechanical stress (By similarity). Required for embryonic development and involved in regulation of several signaling pathways such as MAPK and AKT (By similarity). Required for myoblast differentiation involving regulation of ERK signaling (By similarity). Essential for cardiac homeostasis and proper heart function (By similarity).

Cellular Location
Nucleus inner membrane; Multi-pass membrane protein. Nucleus envelope. Cytoplasm, cytoskeleton, spindle. Note=Lamina-associated protein residing in the inner nuclear membrane (INM) of the nuclear envelope (NE) (PubMed:16339967). The localization to the INM is dependent on LMNA (PubMed:16339967). Evenly distributed around the NE during interphase (PubMed:16339967). During metaphase, found in a reticular network (PubMed:28242692). Recruited to the reforming NE on chromatin disks in early anaphase (PubMed:28242692). In late anaphase, concentrates at the NE core proximal to spindle microtubules, and then broadening to a distributed nuclear rim pattern (PubMed:28242692, PubMed:32494070)

Tissue Location
Ubiquitously expressed, including bone marrow, brain, kidney, colon, skeletal muscle, thymus, testis and uterus


LEMD2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

LEMD2 Antibody - Images


10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



Western blot analysis of LEMD2 in 293 cell lysate with LEMD2 antibody at (A) 1 and (B) 2 µg/mL.



Immunocytochemistry of LEMD2 in 3T3 cells with LEMD2 antibody at 2.5 µg/mL.



Immunofluorescence of LEMD2 in 3T3 cells with LEMD2 antibody at 20 µg/mL.

LEMD2 Antibody - Background

LEMD2 Antibody: The LEM (lamina-associated polypeptide-emerin-MAN1) domain is a motif shared by a group of lamin-interacting proteins of the inner nuclear membrane (INM) and nucleoplasm. One such protein, LEMD2 (LEM domain-containing protein 2), is a lamina-associated protein in the

INM involved in nuclear structure organization, containing an N-terminal LEM motif, two transmembrane domains, and a MAN1-Src1p C-terminal (MSC) domain. LEMD2 is a truncated paralog of MAN1, lacking the MAN1-specific C-terminal RNA-recognition motif, but like MAN1 is required for myogenic differentiation. Mutations in LEMD2 are thought be among the possible causes of muscle disease.

LEMD2 Antibody - References

Brachner A, Reipert S, Foisner R, et al. LEM2 is a novel MAN1-related inner nuclear membrane protein associated with A-type lamins. J. Cell Sci. 2005; 118:5797-810
Huber MD, Guan T, and Gerace L. Overlapping functions of nuclear envelope proteins NET25 (Lem2) and Emerin in regulation of extracellular signal-regulated kinase signaling in myoblast differentiation. Mol. Cell. Biol. 2009; 29:5718-28.