10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

DCP2 Antibody
Catalog # ASC11553

Specification

DCP2 Antibody - Product Information

Application	**WB, IF**
Primary Accession	Q8IU60
Other Accession	NP_689837, 31542498
Reactivity	**Human, Mouse, Rat**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**IgG**
Calculated MW	**46 kDa KDa**
Application Notes	**DCP2 antibody can be used for detection of DCP2 by Western blot at 1 µg/mL. For immunofluorescence start at 20 µg/mL.**

DCP2 Antibody - Additional Information

Gene ID **167227**

Target/Specificity
DCP2; Multiple isoforms of DCP2 are known to exist.

Reconstitution & Storage
DCP2 antibody can be stored at 4℃ for three months and -20℃, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions
DCP2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

DCP2 Antibody - Protein Information

Name DCP2

Synonyms NUDT20

Function
Decapping metalloenzyme that catalyzes the cleavage of the cap structure on mRNAs (PubMed:12417715, PubMed:12218187, PubMed:12923261, PubMed:21070968, PubMed:28002401, PubMed:31875550). Removes the 7-methyl guanine cap structure from mRNA molecules, yielding a 5'-phosphorylated mRNA fragment and 7m-GDP (PubMed:12486012, PubMed:<a href="http://www.uniprot.org/citations/12923261"

target="_blank">12923261, PubMed:21070968, PubMed:28002401, PubMed:31875550). Necessary for the degradation of mRNAs, both in normal mRNA turnover and in nonsense-mediated mRNA decay (PubMed:14527413). Plays a role in replication-dependent histone mRNA degradation (PubMed:18172165). Has higher activity towards mRNAs that lack a poly(A) tail (PubMed:21070968). Has no activity towards a cap structure lacking an RNA moiety (PubMed:21070968). The presence of a N(6)-methyladenosine methylation at the second transcribed position of mRNAs (N(6),2'-O- dimethyladenosine cap; m6A(m)) provides resistance to DCP2-mediated decapping (PubMed:28002401). Blocks autophagy in nutrient-rich conditions by repressing the expression of ATG-related genes through degradation of their transcripts (PubMed:26098573).

Cellular Location
Cytoplasm, P-body. Nucleus Note=Predominantly cytoplasmic, in processing bodies (PB) (PubMed:15273322). A minor amount is nuclear (PubMed:15273322)

Tissue Location
Expressed in brain and testis. Not detected in heart (at protein level).


DCP2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

DCP2 Antibody - Images

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



Western blot analysis of DCP2 in 293 cell lysate with DCP2 antibody at 1 µg/mL in (A) the absence and (B) the presence of blocking peptide.



Immunofluorescence of DCP2 in human testis tissue with DCP2 antibody at 20 µg/mL.

DCP2 Antibody - Background

DCP2 Antibody: The removal, or decapping, of eukaryotic mRNA is an important step in the degradation of mRNA. Decapping protein 2 (DCP2) is the major mRNA decapping enzyme in cells. It is a member of the Nudix hydrolases superfamily of proteins that predominantly catalyze the hydrolysis of small nucleoside diphosphate substrates linked to another moiety. DCP2 is widely expressed in multiple tissues at varying levels, with highest expression seen in testis and brain.

DCP2 Antibody - References

Song MG, Li Y, and Kiledjian M. Multiple mRNA decapping enzymes in mammalian cells. Mol. Cell 2010; 40:423-32.
Dunckley T and Parker R. The DCP2 protein is required for mRNA decapping in Saccharomyces cerevisiae and contains a functional MutT motif. EMBO J. 18:5411-22.
Bessman MJ, Frick DN, and O'Handley SF. The MutT proteins of "Nudix" hydrolases, a family of versatile, widely distributed, "housecleaning" enzymes. J. Biol. Chem. 1996; 271:25059-62