

Pellino 1 Antibody
Catalog # ASC11563**Specification****Pellino 1 Antibody - Product Information**

Application	WB, ICC, IF
Primary Accession	Q96FA3
Other Accession	NP_065702 , 11037063
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	46 kDa KDa
Application Notes	Pellino 1 antibody can be used for detection of Pellino 1 by Western blot at 1 - 2 µg/mL.

Pellino 1 Antibody - Additional Information

Gene ID	57162
Target/Specificity	
PELI1;	

Reconstitution & Storage

Pellino 1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

Pellino 1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Pellino 1 Antibody - Protein Information

Name PELI1

Synonyms PRISM

Function

E3 ubiquitin ligase catalyzing the covalent attachment of ubiquitin moieties onto substrate proteins. Involved in the TLR and IL- 1 signaling pathways via interaction with the complex containing IRAK kinases and TRAF6. Mediates 'Lys-63'-linked polyubiquitination of IRAK1 allowing subsequent NF-kappa-B activation (PubMed:12496252, PubMed:17675297). Mediates 'Lys-48'-linked polyubiquitination of RIPK3 leading to its subsequent proteasome-dependent degradation; preferentially recognizes and mediates the degradation of the 'Thr-182' phosphorylated form of RIPK3 (PubMed:29883609). Negatively regulates necroptosis by reducing RIPK3 expression (PubMed:29883609).

href="http://www.uniprot.org/citations/29883609" target="_blank">29883609). Mediates 'Lys-63'-linked ubiquitination of RIPK1 (PubMed:29883609).

Tissue Location


Expressed at high levels in normal skin but decreased in keratinocytes from toxic epidermal necrolysis (TEN) patients (at protein level).

Pellino 1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Pellino 1 Antibody - Images



Western blot analysis of Pellino 1 in human liver tissue lysate with Pellino 1 antibody at (A) 1 and (B) 2 µg/mL.



Immunocytochemistry of Pellino in HepG2 cells with Pellino 1 antibody at 2.5 µg/ml.



Immunofluorescence of Pellino 1 in HepG2 cells with Pellino 1 antibody at 20 µg/ml.

Pellino 1 Antibody - Background

Pellino 1 Antibody: The Pellino proteins are a highly homologous family of E3 ubiquitin ligases that act as upstream mediators in Toll-like receptor (TLR) pathways that lead to activation of MAP kinases and transcription factors. Pellino 1 is required for interleukin-1-mediated signaling through its interaction with the IRAK4-IRAK-TRAF6 complex, ultimately resulting in the activation of NF-κB. Like other members of the Pellino family, Pellino 1 is an E3 ubiquitin ligase, able to catalyze the polyubiquitination of IRAK1. It is activated via phosphorylation by either IRAK1 and IRAK4 or the IKK-related kinases IKK-ε and TBK1. In addition to phosphorylation, Pellino 1 activity is also modulated via ubiquitination and sumoylation.

Pellino 1 Antibody - References

Moynagh PN. The Pellino family: IRAK E3 ligases with emerging roles in innate immune signaling. *Trends Immunol.* 2009; 30:33-42.
Jiang Z, Johnson J, Nie H, et al. Pellino 1 is required for interleukin-1 (IL-1)-mediated signaling through its interaction with the IL-1 receptor-associated kinase 4 (IRAK4)-IRAK-tumor necrosis factor receptor-associated factor 6 (TRAF6) complex. *J. Biol. Chem.* 2003; 278:10952-6
Butler MP, Hanly JA, and Moynagh PN. Kinase-active interleukin-1 receptor-associated kinases promote polyubiquitination and degradation of the Pellino family: direct evidence for Pellino

proteins being ubiquitin-protein isopeptide ligases. J. Biol. Chem. 2007; 282:29729-37.
Goh ET, Arthur JS, Cheung PC, et al. Identification of the protein kinases that activate the E3
ubiquitin ligase Pellino 1 in the innate immune system. Biochem. J. 2012; 441:339-46