

FKBP12 Antibody

Catalog # ASC11798

Specification

FKBP12 Antibody - Product Information

Application WB, ICC, IF Primary Accession P62942

Other Accession
Reactivity
Host
Rabbit

Clonality Polyclonal Isotype IgG

Calculated MW Predicted: 12 kDa

Observed: 12 kDa KDa

Application Notes

FKBP12 antibody can be used for detection of FKBP12 by Western blot at 1 - 2 µg/ml.

Antibody can also be used for

Immunocytochemistry at 5 μ g/mL. For Immunoflorescence start at 20 μ g/mL.

FKBP12 Antibody - Additional Information

Gene ID 2280

Target/Specificity

FKBP1A; FKBP12 antibody is human, mouse and rat reactive.

Reconstitution & Storage

FKBP12 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

Precautions

FKBP12 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

FKBP12 Antibody - Protein Information

Name FKBP1A

Synonyms FKBP1, FKBP12

Function

Keeps in an inactive conformation TGFBR1, the TGF-beta type I serine/threonine kinase receptor, preventing TGF-beta receptor activation in absence of ligand. Recruits SMAD7 to ACVR1B which prevents the association of SMAD2 and SMAD3 with the activin receptor complex, thereby blocking the activin signal. May modulate the RYR1 calcium channel activity. PPlases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides.

Cellular Location

Cytoplasm, cytosol. Sarcoplasmic reticulum membrane {ECO:0000250|UniProtKB:P62943};




Peripheral membrane protein {ECO:0000250|UniProtKB:P62943}; Cytoplasmic side {ECO:0000250|UniProtKB:P62943}

FKBP12 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

FKBP12 Antibody - Images




Western blot analysis of FKBP12 in A431 cell lysate with FKBP12 antibody at 1 µg/ml.





Immunocytochemistry of FKBP12 in A431 cells with FKBP12 antibody at 5 µg/mL.



Immunofluorescence of FKBP12 in A431 cells with FKBP12 antibody at 20 μg/mL.

FKBP12 Antibody - Background

FKBP, or FK506 binding protein, is a family of proteins that have prolyl isomerase activity and are related to the cyclophilins in function (1). FKBP12, also called FKBP1, was originally identified as a target of FK506 and rapamycin activity and catalyzes the transition between cis and transproline residues critical for proper folding of proteins (2). FKBP12 is an abundant, evolutionarily conserved cytoplasmic protein that is an important regulator of diverse array of cellular processes including T cell activation, entry into the cell cycle and intracellular calcium release (3-5).

FKBP12 Antibody - References

Siekierka JJ, Hung SH, Poe M, et al. A cytosolic binding protein for the immunosuppressant FK506 has peptidyl-prolyl isomerase activity but is distinct from cyclophilin. Nature 1989; 341:755-7. Michnick SW, Rosen MK, Wandless TJ, et al. Solution structure of FKBP, a rotamase enzyme and receptor for FK506 and rapamycin. Science 1991; 252:836-9.

Wang T, Donahoe PK, and Zervos AS. Specific interaction of type I receptors of the TGF-beta family with the immunophilin FKBP-12. Science 1994; 265:674-6.

Wen H, Kang S, Song Y, et al. Characterization of the binding sites for the interactions between FKBP12 and intracellular calcium release channels. Arch. Biochem. Biophys. 2012; 517:37-42.