10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

KIRREL2 Antibody
Catalog # ASC11854

Specification

KIRREL2 Antibody - Product Information

Application	**WB, IHC, IF**
Primary Accession	Q6UWL6
Other Accession	NP_954649, 145275174
Reactivity	**Human, Mouse, Rat**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**IgG**
Calculated MW	**Predicted: 64, 70, 78 kDa** **Observed: 70 kDa KDa**
Application Notes	**KIRREL2 antibody can be used for detection of KIRREL2 by Western blot at 1 - 2 µg/ml. Antibody can also be used for immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL.**

KIRREL2 Antibody - Additional Information

Gene ID **84063**

Target/Specificity
KIRREL2; KIRREL2 antibody is human, mouse and rat reactive. At least five isoforms are are known to exist. This antibody is predicted to not cross-react with other members of the KIRREL protein family.

Reconstitution & Storage
KIRREL2 antibody can be stored at 4℃ for three months and -20℃, stable for up to one year.

Precautions
KIRREL2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

KIRREL2 Antibody - Protein Information

Name KIRREL2

Synonyms NEPH3

Function
May regulate basal insulin secretion.

Cellular Location
Cell membrane {ECO:0000250|UniProtKB:Q7TSU7}; Single-pass type I membrane protein. Note=Localized along the sites of the cell contacts. Colocalizes with E-Cadherin and beta- catenin.

{ECO:0000250|UniProtKB:Q7TSU7}


Tissue Location
Highly expressed in beta-cells of the pancreatic islets.

KIRREL2 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture


KIRREL2 Antibody - Images



Western blot analysis of KIRREL2 in 293 cell lysate with KIRREL2 antibody at 1 µg/ml in (A) the absence and (B) the presence of blocking peptide.



Immunohistochemistry of KIRREL2 in human pancreas tissue with KIRREL2 antibody at 5 µg/ml.



Immunofluorescence of KIRREL2 in human pancreas tissue with KIRREL2 antibody at 20 µg/ml.

KIRREL2 Antibody - Background

KIRREL2, also known as nephrin-like 3, is a type I transmembrane protein belonging to the immunoglobulin superfamily of cell adhesion molecules (1). KIRREL2 consists of five extracellular Ig-like repeats, a transmembrane domain, several glycosylation sites and a cytoplasmic domain that has a stretch of nine conserved residues (1,2). KIRREL2 has been found to be expressed in beta cells of pancreatic islets and is involved in the regulation of both glomerular and neural development (3,4). Downregulation of KIRREL2 is implicated in several proteinuric diseases (4,5).

KIRREL2 Antibody - References

Sun C, Kilburn D, Lukashin A, et al. KIRREL2, a novel immunoglobulin superfamily gene expressed primarily in beta cells of the pancreatic islets. Genomics 2003; 82:130-42.
Heikkila E, Ristola M, Havana M, et al. Trans-interaction of nephrin and Neph1/Neph3 induces cell adhesion that associates with decreased tyrosine phosphorylation of nephrin. Biochem. J. 2011; 435:619-28.
Wang H, Lehtonen S, Chen YC, et al. Neph3 associates with regulation of glomerular and neural development in zebrafish. Differentiation 2012; 83:38-46.
Mizuhara E, Minaki Y, Nakatani T, et al. Purkinje cells originate from cerebellar ventricular zone

progenitors positive for Neph3 and E-cadherin. Dev. Biol. 2010; 338:202-14.