

Beclin 2 Antibody
Catalog # ASC11861**Specification**

Beclin 2 Antibody - Product Information

Application	WB, IHC, IF
Primary Accession	A8MW95
Other Accession	NP_001277622 , 595582373
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 47 kDa

Application Notes	Observed: 55 kDa KDa Beclin 2 antibody can be used for detection of Beclin 2 by Western blot at 1 - 2 µg/ml. Antibody can also be used for immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL.
-------------------	---

Beclin 2 Antibody - Additional Information

Gene ID **441925**

Target/Specificity

BECN1P1; Beclin 2 antibody is human specific. At least two isoforms of Beclin 2 are known to exist; this antibody will only detect the longer isoform. Beclin 2 antibody is predicted to not cross-react with Beclin 1.

Reconstitution & Storage

Beclin 2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

Precautions

Beclin 2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Beclin 2 Antibody - Protein Information

Name BECN2 {ECO:0000303|PubMed:23954414, ECO:0000312|HGNC:HGNC:38606}

Function

Involved in 2 distinct lysosomal degradation pathways: acts as a regulator of autophagy and as a regulator of G-protein coupled receptors turnover. Regulates degradation in lysosomes of a variety of G-protein coupled receptors via its interaction with GPRASP1/GASP1.

Cellular Location

Cytoplasm.

Tissue Location


Present in fetal and adult brain (at protein level).

Beclin 2 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Beclin 2 Antibody - Images



Western blot analysis of Beclin 2 in Jurkat cell lysate with Beclin 2 antibody at 1 μ g/ml in (A) the absence and (B) the presence of blocking peptide.



Immunohistochemistry of Beclin 2 in human lung carcinoma tissue with Beclin 2 antibody at 5 μ g/ml.



Immunofluorescence of Beclin 2 in human lung carcinoma tissue with Beclin 2 antibody at 20 μ g/ml.

Beclin 2 Antibody - Background

Autophagy, the process of bulk degradation of cellular proteins through an autophagosomic-lysosomal pathway is important for normal growth control and may be defective in tumor cells (1,2). Beclin 2 is a mammalian specific homolog of the autophagy protein Beclin 1 (3). Like Beclin 1, Beclin 2 interacts with Bcl-2 and class III PI3K complex components. However, Beclin 2 functions in an additional lysosomal degradation pathway and is required for ligand-induced endolysosomal degradation of several G protein-coupled receptors (3). Beclin 2 is also required for agonist-induced lysosome-mediated degradation of EGFR in lung cancer cells, suggesting that it may also play a role in regulating other intracellular signaling pathways (4).

Beclin 2 Antibody - References

Gozuacik D and Kimchi A. Autophagy as a cell death and tumor suppressor mechanism. *Oncogene*. 2004; 23:2891-906.
Kisen GO, Tessitore L, Costelli P, et al. Reduced autophagic activity in primary rat hepatocellular carcinoma and ascites hepatoma cells. *Carcinogenesis* 1993; 14:2501-5.
He C, Wei Y, Sun K, et al. Beclin 2 functions in autophagy, degradation of G protein-coupled receptors, and metabolism. *Cell* 2013; 154:1085-99.
Zhang W and He C. Regulation of plasma membrane receptors by a new autophagy-related BECN/Beclin family member. *Autophagy* 2014; 10: epub.