

**SPIB Antibody**  
**Catalog # ASC11917****Specification**

---

**SPIB Antibody - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">Q01892</a>
Other Accession	<a href="#">NP_003112</a> , <a href="#">61888836</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 19, 25, 29 kDa; Observed: 20 kDa
Application Notes	SPIB antibody can be used for detection of SPIB by Western blot at 1 - 2 µg/ml. Antibody can also be used for immunohistochemistry starting at 2.5 µg/mL.

**SPIB Antibody - Additional Information**Gene ID **6689****Target/Specificity**

SPIB; SPIB antibody is human, mouse and rat reactive. At least four isoforms of SPIB are known to exist; this antibody will detect all but isoform 3.

**Reconstitution & Storage**

SPIB antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

**Precautions**

SPIB Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**SPIB Antibody - Protein Information****Name** SPIB**Function**

Sequence specific transcriptional activator which binds to the PU-box, a purine-rich DNA sequence (5'-GAGGAA-3') that can act as a lymphoid-specific enhancer. Promotes development of plasmacytoid dendritic cells (pDCs), also known as type 2 DC precursors (pre-DC2) or natural interferon (IFN)-producing cells. These cells have the capacity to produce large amounts of interferon and block viral replication. May be required for B-cell receptor (BCR) signaling, which is necessary for normal B-cell development and antigenic stimulation.

**Cellular Location**

[Isoform 1]: Nucleus


**Tissue Location**

Expressed in plasmacytoid dendritic cells (pDCs) and B-cells, not expressed in T-cells or granulocytes. May also be enriched in stem cell populations of the liver


**SPIB Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**SPIB Antibody - Images**

Western blot analysis of SPIB in human prostate tissue lysate with SPIB antibody at 1 µg/ml.



Immunohistochemistry of SPIB in human prostate tissue with SPIB antibody at 2.5 µg/mL.

### **SPIB Antibody - Background**

SPIB is a member of the ETS transcription factor family that influences lymphoid development and activity and binds the consensus DNA site GGA (A/T) through a unique winged helix-turn-helix motif known as the ETS domain (1). SPIB is found in hematopoietic cells such as B cells and plasmacytoid dendritic cells (DC) (2,3). It promotes the development of plasmacytoid dendritic cells (pDCs) or natural interferon (IFN)-producing cells. SPIB may be required for B-cell receptor (BCR) signaling, which is necessary for normal B-cell development and antigenic stimulation (3,4).

### **SPIB Antibody - References**

Ray D, Bosselut R, Ghysdael J, et al. Characterization of Spi-B, a transcription factor related to the putative oncoprotein Spi-1/PU.1. Mol. Cell Biol. 1992; 12:4297-304.

Nagy M, Chapuis B, and Matthes T. Expression of transcription factors Pu.1, Spi-B, Blimp-1, BSAP and oct-2 in normal human plasma cells and in multiple myeloma cells. Br. J. Haematol. 2002; 116:429-35.

Schotte R, Rissoan MC, Bendriss-Vermare N, et al. The transcription factor Spi-B is expressed in plasmacytoid DC precursors and inhibits T-, B-, and NK-cell development. Blood 2003; 101:1015-23.

Schotte R, Nagasawa M, Weijer K, et al. The ETS transcription factor Spi-B is required for human plasmacytoid dendritic cell development. J. Exp. Med. 2004; 200:1503-9.