10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

alpha-Tubulin Antibody [2B11]

Catalog # ASC12043

Specification

alpha-Tubulin Antibody [2B11] - Product Information

Application	**WB**
Other Accession	37492, CAA25855, 7846
Reactivity	**Human, Mouse, Rat, Rabbit, Zebrafish**
Host	**Mouse**
Clonality	**Monoclonal**
Isotype	**IgG**
Application Notes	**alpha-Tubulin antibody can be used for detection of alpha-Tubulin by Western blot at 1 - 2 µg/ml.**

alpha-Tubulin Antibody [2B11] - Additional Information

Other Names
Tubulin alpha-1A, TUBA1A, TUBA3, LIS3

Precautions
alpha-Tubulin Antibody [2B11] is for research use only and not for use in diagnostic or therapeutic procedures.

alpha-Tubulin Antibody [2B11] - Protein Information


alpha-Tubulin Antibody [2B11] - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

alpha-Tubulin Antibody [2B11] - Images

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



Western blot of alpha-Tubulin in 293, A431, A549, Daudi, HeLa, HepG2, Jurkat, K562, MOLT4, 3T3, Raji, Ramos, THP-1, U937, human brain, mouse brain, rat brain, rabbit brain, mouse lung, rat lung, mouse liver, rabbit spleen and zebrafish lysate at 1 µg/mL.

alpha-Tubulin Antibody [2B11] - Background

Alpha-Tubulin belongs to the tubulin superfamily, which is composed of six distinct families. Along with beta-tubulins, alpha-Tubulins are the major components of microtubules. These microtubules are involved in a wide variety of cellular activities ranging from mitosis and transport events to cell movement and the maintenance of cell shape. Alpha- and beta-tubulin dimers are assembled to 13 protofilaments that form a microtubule of 22-nm diameter (reviewed in 1). Tyrosine ligase adds a C-terminal tyrosine to monomeric alpha-Tubulin. Assembled microtubules can again be detyrosinated by a cytoskeleton-associated carboxypeptidase (2). Another post-translational modification of detyrosinated alpha-Tubulin is C-terminal polyglutamylation, which is characteristic of microtubules in neuronal cells and the mitotic spindle (3). Like GAPDH and ?-Actin, this antibody makes an excellent loading control in immunoblots.

alpha-Tubulin Antibody [2B11] - References

McKean PG, Vaughan S, and Gull K. The extended tubulin family. J. Cell Sci. 2001; 114:2723-33.;Barra HA, Arce CA, and Argarana CE. Posttranslational tyrosination/detyrosination of tubulin. Mol. Neurobiol. 1988; 2:133-53.;Fukshima N, Furuta D, Hidaka Y, et al. Post-translational modifcations of tubulin in the nervous system. J. Neurochem. 2009; 109:683-693.;