10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

alpha-Tubulin Antibody
Catalog # ASC12054

Specification

alpha-Tubulin Antibody - Product Information

Application	**WB, IHC, IF**
Primary Accession	Q71U36
Other Accession	37492, CAA25855, 7846
Reactivity	**Human, Mouse, Rat**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**IgG**
Application Notes	**alpha-Tubulin antibody can be used for detection of alpha-Tubulin by Western blot at 1 - 2 µg/ml. Antibody can also be used for Immunohistochemistry starting at 2.5 µg/mL. For immunofluorescence start at 20 µg/mL.**

alpha-Tubulin Antibody - Additional Information

Other Names
Tubulin alpha-1A, TUBA1A, TUBA3, LIS3

Precautions
alpha-Tubulin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

alpha-Tubulin Antibody - Protein Information

Name TUBA1A

Synonyms TUBA3

Function
Tubulin is the major constituent of microtubules, a cylinder consisting of laterally associated linear protofilaments composed of alpha- and beta-tubulin heterodimers. Microtubules grow by the addition of GTP-tubulin dimers to the microtubule end, where a stabilizing cap forms. Below the cap, tubulin dimers are in GDP-bound state, owing to GTPase activity of alpha-tubulin.

Cellular Location
Cytoplasm, cytoskeleton.

Tissue Location
Expressed at a high level in fetal brain.


10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

alpha-Tubulin Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture


alpha-Tubulin Antibody - Images




Western blot analysis of alpha-Tubulin in 293, Daudi, Hela, HepG2, Jurkat, K562, NH3T3, Raji, Ramos, U937, Human brain, Mouse brain, Rat Brain, Rabbit Brain, Rabbit Spleen, Zebrafish, Mouse Liver and Chicken liver lysate at 1 µg/mL.



Immunohistochemistry of alpha-Tubulin in human brain tissue with alpha-Tubulin antibody at 2.5 µg/ml.



Immunofluorescence of alpha-Tubulin in human brain tissue with alpha-Tubulin antibody at 20 µg/mL.



Immunohistochemistry of alpha-Tubulin in human brain tissue with alpha-Tubulin antibody at 5 µg/mL.

alpha-Tubulin Antibody - Background

alpha-Tubulin belongs to the tubulin superfamily, which is composed of six distinct families. Along with beta-tubulins, alpha-Tubulins are the major components of microtubules. These microtubules are involved in a wide variety of cellular activities ranging from mitosis and transport events to cell movement and the maintenance of cell shape. Alpha- and beta-tubulin dimers are assembled to 13 protofilaments that form a microtubule of 22-nm diameter (reviewed in 1). Tyrosine ligase adds a C-terminal tyrosine to monomeric alpha-Tubulin. Assembled microtubules can again be detyrosinated by a cytoskeleton-associated carboxypeptidase (2). Another post-translational modification of detyrosinated alpha-Tubulin is C-terminal polyglutamylation, which is characteristic of microtubules in neuronal cells and the mitotic spindle (3). Like GAPDH and ?-Actin, this antibody makes an excellent loading control in immunoblots.

alpha-Tubulin Antibody - References

McKean PG, Vaughan S, and Gull K. The extended tubulin family. J. Cell Sci. 2001; 114:2723-33.;Barra HA, Arce CA, and Argarana CE. Posttranslational tyrosination/detyrosination of tubulin. Mol. Neurobiol. 1988; 2:133-53.;Fukshima N, Furuta D, Hidaka Y, et al. Post-translational

modifcations of tubulin in the nervous system. J. Neurochem. 2009; 109:683-693.;