

RAT IgG2b isotype control Fluorescein
Monoclonal IgG2b , Fluorescein (FITC)
Catalog # ASR1129**Specification**

RAT IgG2b isotype control Fluorescein - Product Information

Description	RAT IgG2b isotype control Fluorescein conjugated
Conjugate	Fluorescein (FITC)
FP Value	2-8 moles Fluorescein (FITC) per mole of Rat IgG2b
Clonality	Monoclonal
Application	,4,10,
Application Note	ELISA 1:2000-1:20,000;FlowCytometry 1:1000-1:5000
Physical State	Liquid (sterile filtered)
Host Isotype	IgG2b
Species of Origin	Rat
Stabilizer	None
Preservative	0.01% (w/v) Sodium Azide

RAT IgG2b isotype control Fluorescein - Additional Information**Shipping Condition**

Wet Ice

Purity

RAT IgG2b isotype control has been prepared from concentrated cell culture supernatant by immunoaffinity chromatography using protein G. In an Ouchterlony double diffusion assay the material is non-reactive with antisera to rat IgG1, IgG2a, IgG3 , IgM , and IgA. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rat IgG and anti-Rat serum. Light and heavy chain composition has been confirmed.

Storage Condition

Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. DO NOT FREEZE. This product is light sensitive.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

RAT IgG2b isotype control Fluorescein - Protein Information**RAT IgG2b isotype control Fluorescein - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RAT IgG2b isotype control Fluorescein - Images

RAT IgG2b isotype control Fluorescein - Background

RAT IgG2b isotype control is used in flow cytometry, western blot and ELISA and differentiate between immunoglobulin classes and subclasses. Isotype controls allow for the genetic variations or differences in the constant regions of the heavy and light chains. In Rat there are six relevant heavy chain isotypes and two light chain isotypes: heavy chain α - IgA, γ - IgG 1, 2a, 2b, 2c and μ - IgM, light chain κ and λ .