

Fab Anti-Goat IgG (H&L) Secondary Antibody

Donkey Polyclonal, Unconjugated Catalog # ASR1244

Specification

Fab Anti-Goat IgG (H&L) Secondary Antibody - Product Information

Description F(ab) Anti-GOAT IgG (H&L) (DONKEY)

Antibody Host **Donkey** Conjugate Unconjugated

Target Species Goat Clonality **Polyclonal** Application ,1,10,15,

Application Note ELISA 1:20,000-1:100,000; Western Blot

1:2,000-1:10,000; Immunochemistry

1:1,000-1:5,000

Physical State Liquid (sterile filtered)

Host Isotype IgG F(ab) Target Isotype IqG (H&L)

Buffer 0.02 M Potassium Phosphate, 0.15 M

> Sodium Chloride, pH 7.2 Goat IgG whole molecule

Immunogen None

Stabilizer

Preservative 0.01% (w/v) Sodium Azide

Fab Anti-Goat IgG (H&L) Secondary Antibody - Additional Information

Shipping Condition

Wet Ice

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Donkey Serum. No reaction was observed against anti-Pepsin or anti-Donkey IgG F(c).

Storage Condition

Store vial at 4° C prior to opening. This product is stable at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Fab Anti-Goat IgG (H&L) Secondary Antibody - Protein Information





Tel: 858.875.1900 Fax: 858.875.1999

Fab Anti-Goat IgG (H&L) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Fab Anti-Goat IgG (H&L) Secondary Antibody - Images