

#### Fab Anti-Human IgG (H&L) Secondary Antibody Goat Polyclonal, Unconjugated

Catalog # ASR1247

## Specification

# Fab Anti-Human IgG (H&L) Secondary Antibody - Product Information

Description

Host Conjugate Target Species Reactivity Clonality Application Application Note

Physical State Host Isotype Target Isotype Buffer

Immunogen

Stabilizer Preservative F(ab) Anti-HUMAN IgG (H&L) (GOAT) Antibody Goat Unconjugated Human Human Polyclonal ,1,10,15, ELISA 1:40,000;Western Blot 1:2,000-1:10,000;Immunochemistry 1:1,000-1:5,000 Liquid (sterile filtered) lgG F(ab) IgG (H&L) 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Anti-Human IgG whole molecule was produced by repeated immunization with Human IgG whole molecule in goat. None 0.01% (w/v) Sodium Azide

## Fab Anti-Human IgG (H&L) Secondary Antibody - Additional Information

Shipping Condition Wet Ice

#### Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum. No reaction was observed against anti-Pepsin or anti-Goat IgG F(c).

### Storage Condition

Store vial at 4° C prior to opening. This product is stable at 4° C as an undiluted liquid. Dilute only prior to immediate use.

### **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

## Fab Anti-Human IgG (H&L) Secondary Antibody - Protein Information



### Fab Anti-Human IgG (H&L) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

### Fab Anti-Human IgG (H&L) Secondary Antibody - Images

### Fab Anti-Human IgG (H&L) Secondary Antibody - Background

Anti-Human IgG whole molecule antibody generated in goat detects specifically human IgG whole molecule. This secondary antibody anti-Human is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assays, western-blot, immunoprecipitation and more generally immunoassays.