

Anti-Donkey IgG F(c) Secondary Antibody

Goat Polyclonal, Unconjugated Catalog # ASR1376

Specification

Anti-Donkey IgG F(c) Secondary Antibody - Product Information

Description Host Conjugate Target Species Clonality Application Application Note

Physical State Host Isotype Target Isotype Buffer

Immunogen Reconstitution Volume Reconstitution Buffer

Stabilizer Preservative Anti-DONKEY IgG F(c) (GOAT) Antibody Goat Unconjugated Donkey Polyclonal ,1,10,15, ELISA 1:20,000-1:100,000;Western Blot 1:2,000-1:10,000;Immunochemistry 1:1,000-1:5,000 Lyophilized Antiserum laG F(c) 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Donkey IgG F(c) fragment 2.0 mL Restore with deionized water (or equivalent) None 0.01% (w/v) Sodium Azide

Anti-Donkey IgG F(c) Secondary Antibody - Additional Information

Shipping Condition Ambient

Purity

This product was prepared from monospecific antiserum by a delipidation and defibrination. Assay by immunoelectrophoresis resulted in a single precipitin arc against Donkey IgG, Donkey IgG F(c) and Donkey Serum. No reaction was observed against Donkey IgG F(ab')2.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Donkey IgG F(c) Secondary Antibody - Protein Information



Anti-Donkey IgG F(c) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Donkey IgG F(c) Secondary Antibody - Images