

Anti-Goat IgG (H&L) (Rhodamine Conjugated) Secondary Antibody

Donkey Polyclonal, Rhodamine (TRITC) Catalog # ASR1389

Specification

Anti-Goat IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Product Information

Description

Host Conjugate FP Value

Target Species Clonality Application Application Note

Physical State Host Isotype Target Isotype Buffer

Immunogen Reconstitution Volume Reconstitution Buffer

Stabilizer

Preservative

Anti-GOAT IgG (H&L) (DONKEY) Antibody **Rhodamine Conjugated Donkey Rhodamine (TRITC)** 4.8 moles Rhodamine (TRITC) per mole of laG Goat Polyclonal ,3,4, FLISA 1:10,000-1:50,000;IF Microscopy 1:1,000-1:5,000;FlowCytometry 1:500-1:2.500 Lyophilized lgG IgG (H&L) 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Goat IgG whole molecule 1.0 mL Restore with deionized water (or equivalent) 10 mg/mL Bovine Serum Albumin (BSA) -Immunoglobulin and Protease free 0.01% (w/v) Sodium Azide

Anti-Goat IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Additional Information

Shipping Condition Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Donkey Serum, Goat IgG and Goat Serum.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note



This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Goat IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Protein Information

Anti-Goat IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Goat IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Images

Anti-Goat IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.