

Anti-Donkey IgG (H&L) Secondary Antibody

Goat Polyclonal, Unconjugated Catalog # ASR1442

Specification

Anti-Donkey IgG (H&L) Secondary Antibody - Product Information

Description Host Conjugate Target Species Clonality Application Application Note

Physical State Host Isotype Target Isotype Buffer

Immunogen Stabilizer Preservative Anti-DONKEY IgG (H&L) (GOAT) Antibody Goat Unconjugated Donkey **Polyclonal** ,1,10,15, ELISA 1:20,000-1:100,000;Western Blot 1:2,000-1:10,000;Immunochemistry 1:1,000-1:5,000 Liquid (sterile filtered) laG IaG (H&L) 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Donkey IgG whole molecule None 0.01% (w/v) Sodium Azide

Anti-Donkey IgG (H&L) Secondary Antibody - Additional Information

Shipping Condition Wet Ice

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Donkey IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Donkey IgG and Donkey Serum.

Storage Condition

Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Precautions Note This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Donkey IgG (H&L) Secondary Antibody - Protein Information

Anti-Donkey IgG (H&L) Secondary Antibody - Protocols



Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Donkey IgG (H&L) Secondary Antibody - Images