10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Horse IgG F(c) Biotin
Catalog # ASR1475

Specification

Horse IgG F(c) Biotin - Product Information

Description	**HORSE IgG F(c) fragment Biotin conjugated**
Conjugate	**Biotin**
Physical State	**Lyophilized**
Host Isotype	**IgG F(c)**
Buffer	**0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2**
Species of Origin	**Horse**
Reconstitution Volume	**1.0 mL**
Reconstitution Buffer	**Restore with deionized water (or equivalent)**
Stabilizer	**10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free**
Preservative	**0.01% (w/v) Sodium Azide**

Horse IgG F(c) Biotin - Additional Information

Shipping Condition
Ambient

Purity
This product was prepared from normal serum delipidation, salt fractionation, ion exchange chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Horse IgG, anti-Horse IgG F(c) and anti-Horse Serum. No reaction was observed against anti-Horse IgG F(ab')2 or anti-Papain.

Storage Condition
Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note
This product is for research use only and is not intended for therapeutic or diagnostic applications.

Horse IgG F(c) Biotin - Protein Information

Horse IgG F(c) Biotin - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Horse IgG F(c) Biotin - Images