

Anti-Bovine IgG (H&L) Secondary Antibody

Rabbit Polyclonal, Unconjugated Catalog # ASR1546

Specification

Anti-Bovine IgG (H&L) Secondary Antibody - Product Information

Description Anti-BOVINE IgG (H&L) (RABBIT) Antibody

Host Rabbit

Conjugate
Target Species
Clonality

Unconjugated
Bovine
Polyclonal

Application ,1,2,10,

Application Note ELISA 1:20,000-1:100,000; Western Blot

1:2,000-1:10,000;Immunohistochemistry

1:1,000-1:5,000

Physical State Liquid (sterile filtered)

Host Isotype IgG

Target Isotype IgG (H&L)

Buffer 0.01 M Sodium Phosphate, 0.15 M Sodium

Chloride, pH 7.2

Immunogen Bovine IgG whole molecule

Stabilizer None

Preservative 0.01% (w/v) Sodium Azide

Anti-Bovine IgG (H&L) Secondary Antibody - Additional Information

Shipping Condition

Wet Ice

Purity

Anti-Bovine IgG (H&L) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Bovine IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Bovine IgG and Bovine Serum.

Storage Condition


Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Bovine IgG (H&L) Secondary Antibody - Protein Information

Anti-Bovine IgG (H&L) Secondary Antibody - Protocols





Tel. 000.070.1900 Fax. 000.070.1999

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Bovine IgG (H&L) Secondary Antibody - Images

Anti-Bovine IgG (H&L) Secondary Antibody - Background

Anti-Bovine IgG whole molecule antibody generated in goat detects specifically human IgG whole molecule. This secondary antibody anti-Bovine is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assays, western-blot, immunoprecipitation and more generally immunoassays. Anti-Bovine IgG (H&L) antibody is ideal for investigators in Immunology, Cancer, and Microbology research.