

Anti-Chicken IgG (H&L) (Biotin Conjugated) Secondary Antibody

Goat Polyclonal, Biotin Catalog # ASR1554

Specification

Physical State

Host Isotype

Anti-Chicken IgG (H&L) (Biotin Conjugated) Secondary Antibody - Product Information

Description Anti-CHICKEN IgG (GOAT) Antibody Biotin

Conjugated

Host Goat Conjugate Biotin

FP Value 10-20 moles Biotin per mole of IgG

Target Species Chicken
Clonality Polyclonal
Application ,1,10,15,

Application Note ELISA 1:800,000; Western Blot

1:3,000-1:30,000;Immunochemistry

1:500-1:3,000 Lyophilized

IqG

Target Isotype IgG (H&L)

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2 Chicken IgG whole molecule

Immunogen Chicker
Reconstitution Volume 1.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

Anti-Chicken IgG (H&L) (Biotin Conjugated) Secondary Antibody - Additional Information

Shipping Condition

Ambient

Purity

Anti-Chicken IgG antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Chicken IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Chicken IgG and Chicken Serum.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.



Anti-Chicken IgG (H&L) (Biotin Conjugated) Secondary Antibody - Protein Information

Anti-Chicken IgG (H&L) (Biotin Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Chicken IgG (H&L) (Biotin Conjugated) Secondary Antibody - Images

Anti-Chicken IgG (H&L) (Biotin Conjugated) Secondary Antibody - Background

Chicken IgG antibody detects the heavy and light chains of the chicken Immunogolbulin G molecule. Anti-Chicken IgG (H&L) antibody is ideal for investigators involved in serum protein component research.