

# Anti-GOAT IgG (H&L) (CHICKEN) Secondary Antibody

Chicken Polyclonal, Unconjugated Catalog # ASR1570

## **Specification**

## Anti-GOAT IgG (H&L) (CHICKEN) Secondary Antibody - Product Information

Description Anti-GOAT IgG (H&L) (CHICKEN) Antibody

Host Chicken Conjugate Unconjugated

Target Species
Clonality
Application

Goat
Polyclonal
1,10,15,

Application Note ELISA 1:170,000; Western Blot

1:2,000-1:10,000;Immunochemistry

1:1,000-1:5,000

Physical State Liquid (sterile filtered)

Host Isotype IgG

Target Isotype IgG (H&L)

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2

Immunogen Goat IgG whole molecule

Stabilizer None

Preservative 0.01% (w/v) Sodium Azide

# Anti-GOAT IgG (H&L) (CHICKEN) Secondary Antibody - Additional Information

### **Shipping Condition**

Wet Ice

#### **Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Chicken Serum, Goat IgG and Goat Serum.

### **Storage Condition**

Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.


#### **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

### Anti-GOAT IgG (H&L) (CHICKEN) Secondary Antibody - Protein Information

# Anti-GOAT IgG (H&L) (CHICKEN) Secondary Antibody - Protocols







Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-GOAT IgG (H&L) (CHICKEN) Secondary Antibody - Images