

Anti-Mouse IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody

Goat Polyclonal, Fluorescein (FITC) Catalog # ASR1615

Specification

Anti-Mouse IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody - Product Information

Description

Host Conjugate FP Value

Target Species Clonality Application Application Note

Physical State Host Isotype Target Isotype Buffer

Immunogen Reconstitution Volume Reconstitution Buffer

Stabilizer

Preservative

Anti-MOUSE IgG (H&L) (GOAT) Antibody Fluorescein Conjugated (Min X Human **Serum Proteins**) Goat Fluorescein (FITC) 6.6 moles Fluorescein (FITC) per mole of lqG Mouse Polyclonal ,3,4, IF Microscopy 1:500-1:2,500;FlowCytometry 1:1,000-1:5,000 Lyophilized lqG IqG (H&L) 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Mouse IgG whole molecule 1.0 mL Restore with deionized water (or equivalent) 10 mg/mL Bovine Serum Albumin (BSA) -**Immunoglobulin and Protease free** 0.01% (w/v) Sodium Azide

Anti-Mouse IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody - Additional Information

Shipping Condition Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Goat Serum, Mouse IgG and Mouse Serum. No reaction was observed against Human Serum Proteins.

Storage Condition

Store secondary antibody at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at



4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Mouse IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody -Protein Information

Anti-Mouse IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody -Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Mouse IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody -Images

Anti-Mouse IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody -Background

Mouse secondary antibody conjugated to FITC is available in a variety of formats. Anti IgG secondary antibody conjugated is suitable for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. Anti mouse IgG FITC is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.