

Anti-Donkey IgG (H&L) (Rhodamine Conjugated) Secondary Antibody

Goat Polyclonal, Rhodamine (TRITC)
Catalog # ASR1677

Specification

Target Species

Physical State

Target Isotype

Host Isotype

Immunogen

Reconstitution Volume

Reconstitution Buffer

Buffer

Anti-Donkey IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Product Information

Description Anti-DONKEY IgG (H&L) (GOAT) Antibody

Rhodamine Conjugated

Host Goat

Conjugate Rhodamine (TRITC)

FP Value 2.6 moles Rhodamine (TRITC) per mole of

lgG Donkey Polyclonal

Clonality Polye Application ,3,4,

Application Note FLISA 1:10,000-1:50,000;IF Microscopy

1:1,000-1:5,000;FlowCytometry

1:500-1:2,500 Lyophilized

IgG

IgG (H&L)

0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2

Donkey IgG whole molecule

1.0 mL

Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

Anti-Donkey IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Additional Information

Shipping Condition

Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Donkey IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Donkey IgG and Donkey Serum.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.



Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Donkey IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Protein Information

Anti-Donkey IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-Donkey IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Images

Anti-Donkey IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.