

# Anti-Donkey IgG (H&L) (Biotin Conjugated) Secondary Antibody

Goat Polyclonal, Biotin Catalog # ASR1679

## Specification

# Anti-Donkey IgG (H&L) (Biotin Conjugated) Secondary Antibody - Product Information

Description

- Host Conjugate Target Species Clonality Application Application Note
- Physical State Host Isotype Target Isotype Buffer
- Immunogen Species of Origin Reconstitution Volume Reconstitution Buffer

Stabilizer

Preservative

#### Goat **Biotin** Donkey Polyclonal ,1,2,10, ELISA 1:20,000-1:100,000;Western Blot 1:2,000-1:10,000;Immunohistochemistry 1:1,000-1:5,000 Lyophilized laG IgG (H&L) 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Donkey IgG whole molecule goat 1.0 mL Restore with deionized water (or equivalent) 10 mg/mL Bovine Serum Albumin (BSA) -Immunoglobulin and Protease free 0.01% (w/v) Sodium Azide

Anti-DONKEY IgG (H&L) (GOAT) Antibody

**Biotin Conjugated** 

## Anti-Donkey IgG (H&L) (Biotin Conjugated) Secondary Antibody - Additional Information

Shipping Condition Ambient

#### **Purity**

Anti-Donkey IgG (H&L) was prepared from monospecific antiserum by immunoaffinity chromatography using Donkey IgG coupled to agarose. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Donkey IgG and Donkey Serum.

## **Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

## **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.



## Anti-Donkey IgG (H&L) (Biotin Conjugated) Secondary Antibody - Protein Information

## Anti-Donkey IgG (H&L) (Biotin Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

## Anti-Donkey IgG (H&L) (Biotin Conjugated) Secondary Antibody - Images

## Anti-Donkey IgG (H&L) (Biotin Conjugated) Secondary Antibody - Background

Donkey IgG (H&L) antibody recognizes the heavy and light chains of the Donkey IgG molecule. Anti-Donkey IgG antibody is ideal for investigators involved in serum component protein research.