

F(ab')₂ Anti-Rabbit IgG F(ab')₂ Pre-Adsorbed Secondary Antibody
Goat Polyclonal, Unconjugated
Catalog # ASR1696**Specification****F(ab')₂ Anti-Rabbit IgG F(ab')₂ Pre-Adsorbed Secondary Antibody - Product Information**

Description	F(ab')₂ Anti-RABBIT IgG F(ab')₂ (GOAT) Antibody Min X Bv Hs Hu Ms Rt & Sh Serum Proteins
Host	Goat
Conjugate	Unconjugated
Target Species	Rabbit
Clonality	Polyclonal
Application	,1,10,15,
Application Note	ELISA 1:10,000-1:50,000;Western Blot 1:1,000-1:5,000;Immunochemistry 1:500-1:3,000
Physical State	Liquid (sterile filtered)
Host Isotype	IgG F(ab')₂
Target Isotype	IgG F(ab')₂
Buffer	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Rabbit IgG F(ab')₂ fragment
Stabilizer	None
Preservative	0.01% (w/v) Sodium Azide

F(ab')₂ Anti-Rabbit IgG F(ab')₂ Pre-Adsorbed Secondary Antibody - Additional Information**Shipping Condition**

Wet Ice

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Rabbit IgG, Rabbit IgG F(ab')₂ and Rabbit Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Rabbit IgG F(c) or Bovine, Horse, Human, Mouse, Rat and Sheep Serum Proteins.

Storage Condition

Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.


F(ab')₂ Anti-Rabbit IgG F(ab')₂ Pre-Adsorbed Secondary Antibody - Protein Information

F(ab')₂ Anti-Rabbit IgG F(ab')₂ Pre-Adsorbed Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

F(ab')₂ Anti-Rabbit IgG F(ab')₂ Pre-Adsorbed Secondary Antibody - Images



Western Blot of Goat anti-F(ab')₂ Rabbit IgG F(ab')₂ Antibody (Pre-Adsorbed) Lane 1: Rabbit IgG F(ab')₂ Load: 100 ng per lane Primary antibody: F(ab')₂ Rabbit IgG F(ab')₂ Antibody (Pre-Adsorbed) at 1:1000 o/n at 4°C Secondary antibody: DyLight™ 800 Donkey anti-goat at 1:20,000 for 30 min at RT Block: MB-070 for 30 min at RT Predicted/Observed size: 28 kDa, 28 kDa Other band(s): antigen breakdown

F(ab')₂ Anti-Rabbit IgG F(ab')₂ Pre-Adsorbed Secondary Antibody - Background

F(ab')₂ Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab')₂ fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and experimental applications. F(ab')₂ fragments penetrate into tissue samples and show better antigen recognition and signal generation in IHC. F(ab')₂ fragments lack the Fc region and therefore do not bind Fc receptors which effectively lowers background staining. F(ab')₂ Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.