10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Bovine IgG F(c) Secondary Antibody
Rabbit Polyclonal, Unconjugated
Catalog # ASR1751

Specification

Anti-Bovine IgG F(c) Secondary Antibody - Product Information

Description	**Anti-BOVINE IgG F(c) (RABBIT) Antibody**
Host	**Rabbit**
Conjugate	**Unconjugated**
Target Species	**Bovine**
Reactivity	**Bovine**
Clonality	**Polyclonal**
Application	**,1,10,15,**
Application Note	**ELISA 1:20,000-1:100,000;Western Blot 1:2,000-1:10,000;Immunochemistry 1:1,000-1:5,000**
Physical State	**Liquid (sterile filtered)**
Host Isotype	**IgG**
Target Isotype	**IgG F(c)**
Buffer	**0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2**
Immunogen	**Anti-Bovine IgG F(c) was produced by repeated immunization with bovine IgG F(c) fragment in rabbit.**
Stabilizer	**None**
Preservative	**0.01% (w/v) Sodium Azide**

Anti-Bovine IgG F(c) Secondary Antibody - Additional Information

Shipping Condition
Wet Ice

Purity
This product was prepared from monospecific antiserum by immunoaffinity chromatography using Bovine IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Bovine IgG, Bovine IgG F(c) and Bovine Serum. No reaction was observed against Bovine IgG F(ab')2.

Storage Condition
Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Precautions Note
This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Bovine IgG F(c) Secondary Antibody - Protein Information

Anti-Bovine IgG F(c) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Bovine IgG F(c) Secondary Antibody - Images

Anti-Bovine IgG F(c) Secondary Antibody - Background

Anti-Bovine IgG F(c) antibody generated in rabbit detects specifically bovine IgG. This secondary antibody anti-Bovine is ideal for investigators who routinely perform titration assays, western-blot, immunoprecipitation and more generally immunoassays.