

Anti-Cat IgG F(c) Secondary Antibody

Goat Polyclonal, Unconjugated Catalog # ASR1754

Specification

Anti-Cat IgG F(c) Secondary Antibody - Product Information

Description Host Conjugate Target Species Clonality Application Application Note

Physical State Host Isotype Target Isotype Buffer

Immunogen Stabilizer Preservative Anti-CAT IgG F(c) (GOAT) Antibody Goat Unconjugated Cat Polyclonal ,1,2,10, ELISA 1:20,000-1:100,000;Western Blot 1:2,000-1:10,000;Immunohistochemistry 1:1,000-1:5,000 Liquid (sterile filtered) laG laG F(c) 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Cat IgG F(c) fragment None 0.01% (w/v) Sodium Azide

Anti-Cat IgG F(c) Secondary Antibody - Additional Information

Shipping Condition Wet Ice

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Cat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Cat IgG, Cat IgG F(c) and Cat Serum. No reaction was observed against Cat IgG F(ab')2.

Storage Condition

Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Precautions Note This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Cat IgG F(c) Secondary Antibody - Protein Information

Anti-Cat IgG F(c) Secondary Antibody - Protocols



Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Cat IgG F(c) Secondary Antibody - Images