10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Rabbit IgG F(c) (Fluorescein Conjugated) Secondary Antibody
Goat Polyclonal, Fluorescein (FITC)
Catalog # ASR1816

Specification

Anti-Rabbit IgG F(c) (Fluorescein Conjugated) Secondary Antibody - Product Information

Description	**Anti-RABBIT IgG F(c) (GOAT) Antibody Fluorescein Conjugated**
Host	**Goat**
Conjugate	**Fluorescein (FITC)**
FP Value	**3.5 moles Fluorescein (FITC) per mole of IgG**
Target Species	**Rabbit**
Clonality	**Polyclonal**
Application	**,3,4,**
Application Note	**FLISA 1:10,000-1:50,000;IF Microscopy 1:1,000-1:5,000;FlowCytometry 1:500-1:2,500**
Physical State	**Lyophilized**
Host Isotype	**IgG**
Target Isotype	**IgG F(c)**
Buffer	**0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2**
Immunogen	**Rabbit IgG F(c) fragment**
Reconstitution Volume	**1.0 mL**
Reconstitution Buffer	**Restore with deionized water (or equivalent)**
Stabilizer	**10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free**
Preservative	**0.01% (w/v) Thimerosal**

Anti-Rabbit IgG F(c) (Fluorescein Conjugated) Secondary Antibody - Additional Information

Shipping Condition
Ambient

Purity
This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Goat Serum, Rabbit IgG, Rabbit IgG F(c) and Rabbit Serum. No reaction was observed against Rabbit IgG F(ab')2.

Storage Condition
Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Precautions Note
This product is for research use only and is not intended for therapeutic or diagnostic applications.


Anti-Rabbit IgG F(c) (Fluorescein Conjugated) Secondary Antibody - Protein Information

Anti-Rabbit IgG F(c) (Fluorescein Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Rabbit IgG F(c) (Fluorescein Conjugated) Secondary Antibody - Images



Western Blot of Anti-Rabbit IgG F(c) (GOAT) Antibody . Lane M: 3 µl Molecular Ladder. Lane 1: Rabbit IgG whole molecule . Lane 2: Rabbit IgG F(ab) Fragment . Lane 3: Rabbit IgG F(c) Fragment . Lane 4: Rabbit IgM Whole Molecule . Lane 5: Normal Rabbit Serum . All samples were reduced. Load: 50 ng of IgG, F(ab), IgM and Serum, 100 ng of F(c). Block: MB-070 for 30 min at RT. Primary Antibody: Anti-Rabbit IgG F(c) (GOAT) Antibody 1:2,000 for 60 min at RT. Secondary antibody: Anti-Goat IgG (DONKEY) Peroxidase Conjugated Antibody 1:40,000 in MB-070 for 30 min at RT. Predicted/Obsevered Size: 25 and 50 kDa for Rabbit IgG and Serum, 25 kDa for F(c) and F(ab), 70 and 23 kDa for IgM. Rabbit F(c) migrates slightly higher.

Anti-Rabbit IgG F(c) (Fluorescein Conjugated) Secondary Antibody - Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.