

# Anti-Cat IgG (H&L) (Peroxidase Conjugated) Secondary Antibody

Rabbit Polyclonal, Peroxidase (Horseradish)
Catalog # ASR1882

### **Specification**

### Anti-Cat IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Product Information

Description Anti-CAT IgG (H&L) (RABBIT) Antibody

**Peroxidase Conjugated** 

Host Rabbit

Conjugate Peroxidase (Horseradish)

Target Species Cat

Clonality Polyclonal Application ,1,10,15,

Application Note ELISA 1:90,000; Western Blot

1:5,000-1:20,000;Immunochemistry

1:2,000-1:7,500 Lyophilized

Physical State
Host Isotype
IgG

Target Isotype IgG (H&L)

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2 Cat IgG whole molecule

Reconstitution Volume 1.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Gentamicin Sulfate. Do NOT

add Sodium Azide!

### Anti-Cat IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Additional Information

#### **Shipping Condition**

**Ambient** 

Immunogen

#### **Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Cat IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Cat IgG and Cat Serum.

### **Storage Condition**

Store vial at  $4^{\circ}$  C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at  $4^{\circ}$  C as an undiluted liquid. Dilute only prior to immediate use.

#### **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.



## Anti-Cat IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Protein Information

# Anti-Cat IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Cat IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Images