10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

F(ab')2 Anti-Mouse IgG F(c) Pre-Adsorbed Secondary Antibody

Goat Polyclonal, Unconjugated
Catalog # ASR1971

Specification

F(ab')2 Anti-Mouse IgG F(c) Pre-Adsorbed Secondary Antibody - Product Information

Description	F(ab')2 Anti-MOUSE IgG F(c) (GOAT) Antibody Min X Bv Hs & Hu Serum Proteins
Host	Goat
Conjugate	Unconjugated
Target Species	Mouse
Clonality	Polyclonal
Application	,1,10,15,
Application Note	ELISA 1:10,000;Western Blot 1:2,000-1:10,000;Immunochemistry 1:1,000-1:5,000
Physical State	Liquid (sterile filtered)
Host Isotype	IgG F(ab')2
Target Isotype	IgG F(c)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Mouse IgG F(c) fragment
Stabilizer	None
Preservative	0.01% (w/v) Sodium Azide

F(ab')2 Anti-Mouse IgG F(c) Pre-Adsorbed Secondary Antibody - Additional Information

Shipping Condition
Wet Ice

Purity
This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Mouse IgG, Mouse IgG F(c) and Mouse Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Mouse IgG F(ab')2 or Bovine, Horse and Human Serum Proteins.

Storage Condition
Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Precautions Note
This product is for research use only and is not intended for therapeutic or diagnostic applications.

F(ab')2 Anti-Mouse IgG F(c) Pre-Adsorbed Secondary Antibody - Protein Information

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

F(ab')2 Anti-Mouse IgG F(c) Pre-Adsorbed Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

F(ab')2 Anti-Mouse IgG F(c) Pre-Adsorbed Secondary Antibody - Images

F(ab')2 Anti-Mouse IgG F(c) Pre-Adsorbed Secondary Antibody - Background

F(ab')2 Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and experimental applications. F(ab)2 fragments penetrate into tissue samples and show better antigen recognition and signal generation in IHC. F(ab)2 fragments lack the Fc region and therefore do not bind Fc receptors which effectively lowers background staining. F(ab')2 Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.