10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Rat IgM (mu chain) Secondary Antibody
Rabbit Polyclonal, Unconjugated
Catalog # ASR2058

Specification

Anti-Rat IgM (mu chain) Secondary Antibody - Product Information

Description	**Anti-RAT IgM (mu chain) (RABBIT) Antibody**
Host	**Rabbit**
Conjugate	**Unconjugated**
Target Species	**Rat**
Clonality	**Polyclonal**
Application	**,1,10,15,**
Application Note	**ELISA 1:50,000-1:250,000;Western Blot 1:5,000-1:25,000;Immunochemistry 1:1,000-1:5,000**
Physical State	**Liquid (sterile filtered)**
Host Isotype	**IgG**
Target Isotype	**IgM µ chain**
Buffer	**0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2**
Immunogen	**Rat IgM mu heavy chain**
Stabilizer	**None**
Preservative	**0.01% (w/v) Sodium Azide**

Anti-Rat IgM (mu chain) Secondary Antibody - Additional Information

Shipping Condition
Wet Ice

Purity
This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgM coupled to agarose followed by solid phase adsorption(s). Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Rat IgM and Rat Serum. No reaction was observed against other Rat or human heavy or light chain proteins. Specificity was confirmed by ELISA at less than 1% cross reactivity against Rat IgG and Rat IgA.

Storage Condition
Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Precautions Note
This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Rat IgM (mu chain) Secondary Antibody - Protein Information


10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Rat IgM (mu chain) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Rat IgM (mu chain) Secondary Antibody - Images



Western Blot of Anti-Rat IgM (mu chain) (RABBIT) Antibody . Lane M: 3 µl Molecular Ladder. Lane 1: Rat IgG whole molecule . Lane 2: Rat IgG F(c) Fragment . Lane 3: Rat IgG F(ab) Fragment . Lane 4: Rat IgM Whole Molecule . Lane 5: Rat Serum . All samples were reduced. Load: 50 ng per lane. Block: MB-070 for 30 min at RT. Primary Antibody: Anti-Rat IgM (mu chain) (RABBIT) Antibody 1:2,000 for 60 min at RT. Secondary Antibody: Anti-Rabbit IgG (GOAT) Peroxidase Conjugated Antibody 1:40,000 in MB-070 for 30 min at RT. Predicted/Obsevered Size: 25 and 55 kDa for Rat IgG and Serum, 25 kDa for F(c) and F(ab), 78 and 25 kDa for IgM. Rat F(c) migrates slightly higher.